

AL.2.2002-126

C-2

# RANGE PLANT COMMUNITIES AND CARRYING CAPACITY FOR THE UPPER FOOTHILLS SUBREGION OF ALBERTA



**Alberta**

SUSTAINABLE RESOURCE DEVELOPMENT



Introduction

# RANGE PLANT COMMUNITY TYPES AND CARRYING CAPACITY FOR THE UPPER FOOTHILLS SUBREGION

Methods

## Fourth approximation

(This publication is a revision of the second and third approximation of the Range Plant Communities and Carrying Capacity for the Upper Foothills Publication No. T/356)

Results

Native grass and shrublands

Key to grasslands

2001

Key to modified grass and shrublands

Key to native shrublands

Prepared by

A. Grasslands

**Michael G. Willoughby**

UPA1. Water-edge meadow	39
UPA2. Sedge-Slender wheatgrass/Meadow rue	40
UPA3. Tufted hairgrass-Sedge	41
UPA4. Tufted hairgrass-Sedge-Slender wheatgrass	42
UPA5. Rough fescue-Tufted hairgrass	43
UPA6. Rough fescue-Hairy willows	44
UPA7. Rough fescue/Bearberry	45
UPA7a. California oatgrass-Rough fescue/Bearberry	46
UPA8. California oatgrass-Sedge	47
UPA9. Junegrass-Sedge	48
UPA10. Bearberry-Slender wheatgrass	49
UPA11. Fireweed/Hairy willows/Portulacaceae	50
UPA12. Rough fescue-Sage sedge	51
UPA13. Alpine rough fescue	52
UPA14. Cow parsnip-Meadow rue/Pronged thorn	53
UPA15. Hairy willows-Sedge	54
UPA16. Hairy willows-Rough fescue/Bearberry	55
UPA17. Hairy fescue-Parry oatgrass-Sedge	56

Edmonton  
2001

Alberta Sustainable Resource Development  
Public Lands Division

**Pub. no.: T/003**  
**ISBN: 0-7785-1792-6**

For copies of this report contact:

- Michael Willoughby  
9920 108 St., 9th Floor  
Edmonton, Alta.  
T5K 2M4  
(403) 422-4598

E-mail: [mike.willoughby@gov.ab.ca](mailto:mike.willoughby@gov.ab.ca)

## Table of contents

	Page
Introduction	1
Climate of the Upper Foothills subregion	1
Methods	3
How to use this guide	4
Results	5
Native grass and shrublands	31
Key to grasslands	36
Key to modified grass and shrublands	37
Key to native shrublands	38
A. Grasslands	
UFA1. Water sedge meadows	39
UFA2. Sedge-Slender wheatgrass/Meadow rue	40
UFA3. Tufted hairgrass-Sedge	41
UFA4. Tufted hairgrass-Sedge-Slender wheatgrass	42
UFA5. Rough fescue-Tufted hairgrass	43
UFA6. Rough fescue-Hairy wildrye	44
UFA7. Rough fescue/Bearberry	45
UFA7a. California oatgrass-Rough fescue/Bearberry	46
UFA8. California oatgrass-Sedge	47
UFA9. Junegrass/Sedge	48
UFA10. Bearberry/Slender wheatgrass	49
UFA11. Fireweed/Hairy wildrye(Forb meadow)	50
UFA12. Rough fescue-Bog sedge	51
UFA13. Alpine rough fescue	52
UFA14. Cow parsnip-Meadow rue/Fringed brome	53
UFA15. Hairy wildrye-Sedge	54
UFA16. Hairy wildrye-Rough fescue/Bearberry	55
UFA17. Idaho fescue-Parry oatgrass-Sedge	56

## B. Shrublands

UFB1. Willow-Bog birch/Water sedge	57
UFB2. Willow/Slender wheatgrass-Sedge	58
UFB3. Willow/Tufted hairgrass	59
UFB4. Willow/Rough fescue	60
UFB5. Bog birch/Rough fescue/Bearberry	61
UFB6. Willow/California oatgrass-Sedge	62
UFB7. Pussy willow shrubland	63
UFB8. Willow/Hairy wildrye-Sedge	64
UFB9. Bog birch/Sedge-Marsh reedgrass	65
UFB10. Willow-Bog birch/Sedge	66
UFB11. Willow-Bog birch	67
UFB12. Alder-Willow/Horsetail	68

## C. Shrublands and Grasslands modified by grazing 69

UFC1. Slender wheatgrass-Sedge/Low forbs	72
UFC2. Rocky Mtn. fescue/Graceful cinquefoil	73
UFC3. Kentucky bluegrass/Dandelion-Clover	74
UFC4. Kentucky bluegrass-Sedge/Dandelion	75
UFC5. Tufted hairgrass-Kentucky bluegrass	76
UFC6. Sedge-Tufted hairgrass	77
UFC7. Creeping red fescue/Clover	78
UFC8. Kentucky bluegrass-Timothy/Meadow rue	79
UFC9. Purple oatgrass-Rough fescue	80
UFC10. Willow/Kentucky bluegrass	81
UFC11. Sedge-Slender wheatgrass-Rough fescue	82

## D. Deciduous community types 83

### Key to deciduous community types 85

UFD1. Aspen/Rose/Bearberry	87
UFD2. Balsam poplar-White spruce/Willow/Yellow Mtn. avens	88
UFD3. Aspen/Rose/Hairy wildrye	89
UFD4. Aspen/Buffaloberry/Hairy wildrye	90
UFD5. Aspen/Marsh reedgrass	91
UFD6. Balsam poplar/Willow/Horsetail	92

## E. Conifer community types 93

Key to conifer community types	95
UFE1. Lodgepole pine/Bog cranberry	97
UFE2. Lodgepole pine-White spruce/Bunchberry	98
UFE3. Lodgepole pine/Willow/Moss	99
UFE4. Lodgepole pine/Marsh reedgrass	100
UFE5. Black spruce/Willow	101
UFE6. White spruce/Horsetail/Moss	102
UFE7. White spruce/Willow	103
UFE8. White spruce/Bearberry	104
UFE9. White spruce/Juniper-Buffaloberry	105
UFE10. White spruce/Moss	106
F. Cutblock and burn community types	107
UFF1. Juniper/Hairy wildrye	108
UFF2. Rose/Hairy wildrye	109
UFF2a. Fireweed/Hairy wildrye	110
UFF3. White spruce/Horsetail/Kentucky bluegrass	111
UFF4. White spruce/Moss (Cutblock)	112
UFF4a. Lodgepole pine-White spruce/Moss	113
UFF5. River alder-Willow/Fireweed-Cow parsnip	114
UFF6. Aspen/Fireweed	115
UFF7. Aspen/Blueberry-Bearberry/Hairy wildrye	116
UFF8. Kentucky bluegrass-Creeping red fescue/Clover	117
Literature cited	118

### List of Figures

Figure 1. Overview of native shrub and grassland complex in the Upper Foothills subregion	31
Figure 2. Typical native shrub and grassland transition zone in the Upper Foothills subregion	32
Figure 3. Layout of plant community types for native grass and shrublands in the landscape of the Upper Foothills subregion	35
Figure 4. Heavily grazed Kentucky bluegrass/Dandelion community type	69
Figure 5. Aspen/Marsh reedgrass community type on south facing slopes in the Upper Foothills subregion	83
Figure 6. The Lodgepole pine-White spruce/Bunchberry community is the dominant conifer community type in the Upper Foothills	

subregion	93
Figure 7. The Juniper/Hairy wildrye community type results from the harvesting of a White spruce/Juniper community	107

### List of Tables

Table 1. Ecosites and ecosite phases of the Upper Foothills subregion	6
c5 yellow mountain avens	14
c6 hairy wildrye sedge	15
c2 hairy wildrye Aw harvested	16
c4 hairy wildrye Sw harvested	17
e1b tall bilberry/arnica Pl harvested	18
e1b tall bilberry/arnica Sw harvested	19
ff fescue-california oatgrass ecosite	20
ff1 grassland	21
ff2 shrubland	22
ff1 grazed grassland	23
ff2 grazed shrubland	24
f4 bracted honeysuckle harvested	25
g2 forb meadow grazed	26
g3 grass meadow	27
g3 grass meadow grazed	28
j2 horsetail Pb	29
j1 horsetail Sw harvested	30
Table 2. Native grass and shrublands of the Upper Foothills subregion	33
Table 3. Shrubbylands and grasslands modified by heavy grazing in the Upper Foothills subregion	71
Table 4. Deciduous community types within the Upper Foothills subregion	86
Table 5. Conifer community types of the Upper Foothills subregion	96

### List of Maps

Map 1. Location of the Upper Foothills subregion in Alberta	2
---	---

## Abstract

The Upper Foothills subregion is found elevationally below the Subalpine and above the Lower Foothills subregions. It is dominated by closed canopied lodgepole pine forests. In the valley bottoms the shrub and grassland community types are a classic example of multiple use land, providing summer range for livestock, prime habitat for many species of wildlife, productive watersheds, and recreational areas. Despite the importance of these vegetation types for livestock grazing, there is little information available on how grazing affects their production. There is little information on forage productivity, carrying capacity and the associated community types with grazing. The lack of information makes it very difficult to development management prescriptions for multiple use. As a result a "Carrying capacity guide" was developed for the Upper Foothills subregion to provide a framework that would easily group the vegetative community types. It is hoped this classification system can be used by field staff to assess carrying capacity and evaluate range condition on lands within the region.

This guide represents the analysis of 470 plots described in the Upper Foothills subregion, near Grande Cache (Willmore Wilderness Park) and west of Rocky Mtn. House during the summers of 1990-1999. The 470 plots represent 67 community types. These types are split into:

- |                           |                    |
|---------------------------|--------------------|
| A. Native grasslands      | 18 community types |
| B. Native shrublands      | 12 community types |
| C. Grazing modified types | 11 community types |
| D. Deciduous types        | 6 community types  |
| E. Conifer types          | 10 community types |
| F. Cutblocks and burns    | 10 community types |

The dominant plant species, canopy cover, environmental conditions, response to grazing, forage production and carrying capacity are outlined for each type.

Digitized by the Internet Archive  
in 2016

## **Introduction**

The province of Alberta is covered by a broad spectrum of vegetation regions from prairie in the South, to alpine vegetation in the mountains and dense forests in the Central and Northern parts of the province. These broad vegetation regions have been classified into 6 natural regions and 20 subregions for the province (Dept. of Environmental Protection, 1994). Each of the regions consists of groups of plant communities which are influenced by environmental conditions and human impacts. Intensive management of these regions requires the ability to recognize the vegetative communities that have similar productivities and respond to disturbance in the same way. The increase in use of Alberta's northern forests has recently stimulated efforts to develop detailed classification systems. Some of these classification systems include Field guide to Forest ecosystems of West Central Alberta (Corns and Annas, 1986) and Field Guide to Ecosites of West-Central Alberta (Beckingham et al., 1996).

The vegetative communities in the province of Alberta are highly regarded by most resource managers for their ability to provide a wide variety of benefits. They are a classic example of multiple use land, providing summer range for livestock, prime habitat for many species of wildlife, productive watersheds and recreational areas. Despite the importance of these vegetation types for livestock grazing, there is little information available on how grazing affects their production. Specifically, there is little data on the levels of utilization which are detrimental to communities growth. There is also no data on forage productivity, carrying capacity and associated community types with grazing. Traditionally, these community types have been rated at 5 ac/AUM or 60 ac/head/year, but recent work has shown that productivity can vary significantly depending upon the ecological conditions of the site.


The purpose of this guide was to develop a framework that would easily group the vegetative community types in the Upper Foothills subregion of the province. The ultimate goal is a classification system that can be used by the field staff to assess carrying capacity and evaluate range condition on lands within the region. This guide supplements the work done by Beckingham et al. (1996) on the forested community types in the Upper foothills subregion. Their guide describes 65 community types on 13 ecosites. Beckingham's guide is a good description of the forested community types found within the subregion, but it does not include forage production values and carrying capacities. It also only provides a brief description of the native shrubland and grassland communities which are extensively utilized by livestock and wildlife in this subregion.

### **Climate of the Upper Foothills subregion**

This subregion is found elevationally below the subalpine and above the Lower Foothills subregions. It ranges in elevation from 1200-1500m at lower latitudes and from 1000-1250 m at higher latitudes. It is dominated by closed canopy lodgepole pine forests with the potential climax species on reference sites being white spruce and black spruce. This

subregion can be distinguished from the Subalpine subregion by the lack of engelmann spruce and from the Lower Foothills by the lack of aspen.

This subregion has a boreal climate which is modified by the Rocky Mountains. The average annual precipitation is 538 mm with over half the precipitation recieved in the summer months (340 mm). The temperature averages 11.5 °C in the summer and -6.0 °C in the winter. These temperatures are milder and not nearly as extreme as the other subregions within the Boreal forest and Foothills natural regions.



**Map 1. Location of Upper Foothills subregion in Alberta**

## Methods

A community type approach (Mueggler, 1988) to classification was chosen in preference to the habitat type approach (Daubenmire, 1952) or ecosystem association approach (Corns and Annas, 1986) because of the lack of understanding of the successional sequences of the communities. Community types are aggregates of similar plant communities based upon existing floristics regardless of successional status (Mueggler, 1988). Community types are what is actually seen in the field. After defining the community types, they then can be linked to the ecosystem associations developed by Corns and Annas (1986) and Beckingham (1994). In the mean time community types can be used as the basis for mapping and range management planning.

Initially, grazing dispositions were inventoried by forest region in order to develop management plans following the procedure outlined in the Rangeland Resource Information System (1991). Individual plots were initially classified within a forest region using cluster analysis (SAS) and ordination (DECORANA, Gauch, 1982). These types were described in individual carrying capacity guides for each forest. This led to differences in classification of the same types between forests, particularly for deciduous forest types. In an effort to standardize the community name and gain some understanding of each community types ecology, all plots sampled in each forest were reclassified. As the study progressed it became quite evident that there were differences in the productivity of the communities between ecoregions. As a result, it was decided to develop the classification within the ecoregion framework (Strong and Leggat, 1992). An ecoregion is a geographical area that has broad vegetation zones combined with climatic data (Strong and Anderson, 1980). As a result, the vegetation within each ecoregion is strongly influenced by the climatic conditions. Recently, the department has adapted the Natural and Subregions of Alberta classification system. This system incorporates the Natural regions and subregions classification used by Parks with the Ecoregions of Alberta classification used by Forestry, Lands and Wildlife. The Upper Foothills subregion and Upper Boreal Cordilleran ecoregion share the same boundary.

Sampling for this guide occurred within the Upper Foothills subregion. This guide outlines the classification of 439 plots described during the summers of 1990-2000 in the Southern and Northern East Slope Forest regions.

A plot consisted of a 10x10 m macroplot and ten randomly selected 1x1 m microplots to record the canopy cover of shrubs and ten nested 20x50 cm microplots to record the canopy cover of forbs and grass. The inventory followed the Range Survey Manual (1992) and uses the MF5 form. At each macroplot a 50x100 cm was clipped and separated into trees, shrubs, forbs and graminoids, oven dried and weighed. The recommended stocking rate is based on 25 percent of the total production for forested types and 50% total production for grass and shrubland types and the fact that one animal requires 455 kg of dry weight material for one month of grazing.

## How to use the guide

First decide what category the community type is in. If it is in the **Native grass and Shrub category** it will not have tree cover and be found on steep south facing slopes or moist lowland areas adjacent to streams and rivers. The predominant species will be native grasses, willow and bog birch. The **Grazing modified** community types will resemble the native shrub and grassland community types, but will show signs of extensive grazing pressure. These community types will be dominated by grazing resistant species Kentucky bluegrass, clover and dandelion. A couple of moderately grazed community types with a predominant native species cover are also found in this category.

The **Deciduous category** will be plant communities dominated by deciduous tree species aspen and balsam poplar and the **Conifer category** will be plant communities dominated by white spruce, lodgepole pine or black spruce tree species.

In order to understand how the community types in this guide are related to the ecosites and ecosite phases outlined in "Ecosites of West-Central Alberta" (Beckingham et al., 1996), the community types in this guide are arranged by ecosite and ecosite phase (Table 1). **Ecosites** are defined as ecological units that develop under similar environmental influences (climate, moisture and nutrient regime). An **ecosite phase** is a subdivision of the ecosite based on the dominant species in the canopy. Table 1 is a reproduction of Figure 14 in the Ecosites of West-Central Alberta guide with the community types in this guide highlighted. For the most part the ecosites and ecosite phases are the same, particularly for the forested community types, but a number of new ecosites and ecosite phases had to be created for the grass and shrubland community types (Table 1). These included (ff)(mesic/rich) fescue-california oatgrass ecosite, and the (c5) yellow mtn avens, (c6) hairy wildrye grassland, (ff1)grassland, (ff2) shrubland, (g3) grass meadow and (j2) horsetail Pb ecosite phases. The "Grazing succession" and "Harvesting succession" categories (Table 1) outline the successional sequence the community types will undergo with increased grazing pressure or harvesting. A number of new ecosite phases were created for these categories. These included (c2) harvested Aw, (c4) harvested Sw, (e1) tall bilberry/arnica P1-Sw harvested, (ff1) grazed grassland, (ff2) grazed shrubland, (f4)bracted honeysuckle Sw harvested, (g2) grazed forb meadow, (g3) grazed grass meadow and (j1) horsetail Sw harvested. All of the new ecosites and ecosite phases are summarized within this guide.

## Results

The analysis of the 470 plots distinguished 67 community types. These types were split into 6 categories:

- A.) Native grasslands (18 types)
- B.) Native shrublands (12 types)
- C.) Grazing modified types (11 types)
- D.) Deciduous types (6 types)
- E.) Conifer types (10 types)
- F.) Cutblock types (10 types)

The dominant plant species, canopy cover, environmental conditions, forage production and carrying capacity are outlined for each community type.

Table 1. Ecosites, ecosite phases and plant community types for the Upper Foothills subregion (adapted from Beckingham et al., 1996) (Highlighted community types are described in this guide, non-highlighted communities are outlined in the guide Ecosites of West-Central Alberta)

Ecosite	Ecosite phase	Forested community type	Range community type	Grazing succession	Harvesting Succession
a grassland (xeric/poor)	a1 shrubby grassland	a1.1 bearberry grassland	UFA10 Bearberry/Slender wheatgrass		
		a1.2 saskatoon-prickly rose grassland	UFA9 Junegrass-Sedge/Sage		
b bearberry/lichen (subxeric/poor)	b1 bearberry/lichen P1	b1.1 P1/bearberry			
		b1.2 P1/Labrador tea/lichen			
		b1.3 P1/hog cranberry			
c hairy wild rye (submesic/medium)	c1 hairy wild rye P1	c1.1 P1/Canada buffaloberry/hairy wild rye			
		c1.2 P1/green alder/hairy wild rye			
		c1.3 P1/hairy wild rye			
	c2 hairy wild rye Aw	c2.1 Aw/hairy wild rye	UFD1 Aw/Rose/Bearberry UFD3 Aw/Rose/Hairy wildrye		UFF6 Aw/Fireweed UFF7 Aw/Blueberry- Bearberry/Hairy wildrye
c3 hairy wild rye Aw-Sw- P1	c3.1 Aw-Sw-P1/Canada buffaloberry/hairy wild rye	UFD4 Aw/Buffaloberry/ Hairy wildrye			





ff fescue-california oatgrass grassland (mesic/rich)	ff1 grassland	<p>UFA5 Rough fescue-Tufted hairgrass          UFA6 Rough fescue-Hairy wildrye          UFA7 Rough fescue-California oatgrass/Bearberry          UFA7a California oatgrass-Rough fescue/Bearberry          UFA8 California oatgrass-Sedge          UFA12 Rough fescue-Bog sedge          UFA13 Arctic rough fescue</p>	<p>UFC2 Rocky Mtn. fescue/Graceful cinquefoil          UFC7 Creeping red fescue/          Clover          UFC10 Purple oatgrass-Rough fescue          UFC11 Slender wheatgrass-Sedge-Rough fescue          UFA16 Hairy wildrye-Rough fescue/          Bearberry          UFA17 Idaho fescue-Parry oatgrass-Sedge</p>	
	ff2 shrubland	<p>UFB4 Willow/Rough fescue          UFB5 Bog birch/Rough fescue/Bearberry          UFB6 Willow/California oatgrass-Sedge          UFB8 Willow/Hairy wildrye-Sedge</p>	<p>UFC9 Willow/Kentucky bluegrass</p>	
f bracted honeysuckle (Subhygric/rich)	f1 bracted honeysuckle P1	<p>f1.1 P1/green alder/fern          f1.2 P1/bracted honeysuckle/fern          f1.3 P1/fir/fern/feather moss          f1.4 P1/fern/feather moss</p>	<p>UFE3 P1/Willow/Moss</p>	


	f2 bracted honeysuc kle Pb	f2.1 Pb/green alder-river alder/fern															
		f2.2 Pb/bracted honeysuckle/fern				UFD5 Aw/Marsh reedgrass											
	f3 bracted honeysuc kle Pb- Sw-P1	f3.1 Pb-Sw-P1/green alder- river alder/fern															
		f3.2 Pb-Sw-P1/fern/feather moss															
	f4 bracted honeysuc kle Sw	f4.1 Sw/green alder/fern									UFFS River alder-Willow/ Fireweed-Cow parsnip						
		f4.2 Sw/bracted honeysuckle/fern															
		f4.3 Sw/fir/fern/feather moss															
		f4.4 Sw/fern/feather moss															
	f5 bracted honeysuc kle Fa	f5.1 Fa/fir/fern/feather moss															
		f6.1 willow/cow parsnip-fern													UFB12 Alder-Willow/Horsetail		
	f6 bracted honeysuc kle willow	g1.1 willow/cow parsnip-tall larkspur meadow															





m2 shrubby rich fen	m2.1 dwarf birch/sedge/golden moss	UFB1 Willow-Bog birch/Water sedge		
	m2.2 willow/sedge/golden moss			
m3 graminoid rich fen	m3.1 sedge rich fen	UFA1 Water-Beaked sedge meadow		

**c5 Yellow mountain avens (n=1)**



**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
 Nutrient regime: medium, poor  
 Topographic position: crest, upper slope, midslope  
 Slope: level  
 Aspect: northerly, level

**SOIL CHARACTERISTICS**

Organic thickness: (6-15), (0-5)  
 Humus form: mor, moder  
 Surface texture: SiL, L, CL, LS, C  
 Effective texture: SCL, CL, SiL, SL, L, C  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well, rapidly  
 Parent material: M, M/R, GF  
 Soil subgroup: BR.GL, O.EB, E.EB, O.GL

**RANGE PLANT COMMUNITY TYPES**

UFD2. Pb-Sw/ Willow/Yellow Mtn. avens n=1

**CHARACTERISTIC SPECIES**

**Tree**

- [ 7 ] Balsam poplar
- [ 5 ] White spruce

**Shrub**

- [ 16 ] Yellow mountain avens
- [ 13 ] Willow
- [ 9 ] Buffaloberry
- [ 3 ] Bearberry


**Forb**

- [ 11 ] Alpine hedysarum
- [ 11 ] Scouring rush
- [ 4 ] Alpine milk vetch

**Grasses**

- [ 2 ] Blunt Sedge

**c6 Hairy wildrye grassland (n=1)**



**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
 Nutrient regime: medium, poor  
 Topographic position: crest, upper slope, midslope  
 Slope: level  
 Aspect: northerly, level

**SOIL CHARACTERISTICS**

Organic thickness: (6-15), (0-5)  
 Humus form: mor, moder  
 Surface texture: SiL, L, CL, LS, C  
 Effective texture: SCL, CL, SiL, SL, L, C  
 Depth to Mottles: Gley: none  
 Drainage: well, moderately well, rapidly  
 Parent material: M, M/R, GF  
 Soil subgroup: BR.GL, O.EB, E.EB, O.GL

**RANGE PLANT COMMUNITY TYPES**

UFA15. Hairy wildrye-Sedge n=1

**CHARACTERISTIC SPECIES**

**Shrub**

[ 4 ] Bearberry

**Forb**


- [ 5 ] Fireweed
- [ 9 ] Showy locoweed
- [ 4 ] American vetch
- [ 8 ] Strawberry
- [ 2 ] Graceful cinquefoil

**Grasses**

- [ 31 ] Hairy wildrye
- [ 3 ] Arctic bluegrass
- [ 3 ] Sedge
- [ 1 ] Slender wheatgrass

**c2b Harvested hairy wildrye Aw (n=3)**

[ 4 ] Hairy wildrye  
[ 2 ] Indian ricegrass



**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
Nutrient regime: medium, rich  
Topographic position: upper slope, midslope, level  
Slope: (3-5)%  
Aspect: north, east

**SOIL CHARACTERISTICS**

Organic thickness: (6-15), (0-5)  
Humus form: mor  
Surface texture: SiL, L, CL, LS, SiCL, C  
Effective texture: SCL, SiCL, CL, SiC, SL, C  
Depth to Mottles/Gley: none  
Drainage: well, moderately well  
Parent material: M, C, X  
Soil subgroup: BR.GL, O.MB, O.GL, O.EB, O.DYB, GL.GL, E.EB, D.GL

**RANGE PLANT COMMUNITY TYPES**

UFF6. Aw/fireweed n=1  
UFF7. Aw/Blueberry-Bearberry/Hairy wildrye n=2

**CHARACTERISTIC SPECIES**

**Trees**

- [ 1 ] White spruce
- [ 7 ] Aspen
- [ 1 ] Lodgepole pine

**Shrub**

- [ 7 ] Blueberry
- [ 2 ] Bog cranberry
- [ 4 ] Prickly rose
- [ 2 ] Green alder


**Forb**

- [ 18 ] Fireweed
- [ 6 ] Horsetail
- [ 3 ] Heart leaved arnica

**Grasses**

- [ 4 ] Marsh reedgrass
- [ 1 ] Sedge

**c4b Harvested hairy wildrye/ Sw**  
(n=14)



**CHARACTERISTIC SPECIES**

**Tree**

- [10] White spruce
- [7] Balsam poplar
- [3] Aspen

**Understory tree**

- [2] White spruce
- [2] Balsam poplar
- [1] Aspen

**Shrub**

- [3] Shrubby cinquefoil
- [5] Rose
- [7] Creeping juniper
- [8] Willow
- [3] Bearberry

**Forb**

- [3] Showy locoweed
- [3] Northern hedysarum
- [5] Northern bedstraw

**Grasses**

- [3] Sedge
- [21] Hairy wildrye
- [3] Slender wheatgrass

**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
 Nutrient regime: medium, poor  
 Topographic position: upper slope, midslope, level  
 Slope: (2-10)%  
 Aspect: southeasterly, southwesterly, south


**SOIL CHARACTERISTICS**

Organic thickness: (6-15), (0-5)  
 Humus form: mor  
 Surface texture: SiL, L, CL, LS, SiCL, C  
 Effective texture: SCL, SiCL, CL, SiC, SL, C  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: M/R, FL, C/M  
 Soil subgroup: O.EB, E.EB, BR.GL

**RANGE PLANT COMMUNITY TYPES**

- UFF1. Juniper/Hairy wildrye n=4
- UFF2. Rose/Hairy wildrye n=10

**e1b Harvested Pl/ Tall bilberry/  
Feather moss (n=27)**



**SITE CHARACTERISTICS**

Moisture regime: mesic to subxeric  
 Nutrient regime: medium, poor  
 Topographic position: midslope, level, upper slope  
 Slope: (2-41)%  
 Aspect: variable

**SOIL CHARACTERISTICS**

Organic thickness: (0-4)  
 Humus form: moder  
 Surface texture: SiL  
 Effective texture: SiL  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: E, GF  
 Soil subgroup: O.EB

**RANGE PLANT COMMUNITY TYPES**

UFF2a. Fireweed/Hairy wildrye n=22  
 UFF8. Kentucky bluegrass-C. red fescue/Clover n=5

**CHARACTERISTIC SPECIES**

**Shrub**

[ 1 ] Rose

**Forb**

[ 4 ] Fireweed

[ 3 ] Clover

**Grasses**


[ 2 ] Creeping red fescue

[ 3 ] Kentucky bluegrass

[ 8 ] Hairy wildrye

[ 2 ] Timothy

**e3b Harvested tall bilberry/ Arnica/  
Sw (n=6)**



**CHARACTERISTIC SPECIES**

**Tree**

- [ 3 ] White spruce
- [ 10 ] Lodgepole pine

**Shrub**

- [ 1 ] Willow
- [ 2 ] Bunchberry

**Forb**

- [ 1 ] Fireweed
- [ 5 ] Horsetail

**Grasses**

- [ 6 ] Hairy wildrye
- [ 3 ] Marsh reed grass

**Moss**

- [ 1 ] Feather moss
- [ 6 ] Stair step moss

**SITE CHARACTERISTICS**

Moisture regime: mesic to subxeric  
 Nutrient regime: medium, poor  
 Topographic position: midslope, level, upper slope  
 Slope: (2-41)%  
 Aspect: variable

**SOIL CHARACTERISTICS**

Organic thickness: (0-4)  
 Humus form: moder,  
 Surface texture: SiL  
 Effective texture: SiL  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: E, GF  
 Soil subgroup: O, EB

**RANGE PLANT COMMUNITY TYPES**

UFF4a. Pl-Sw/Moss n=6

**ff Fescue-California oatgrass  
grassland (n=70)**

**mesic/rich**

**GENERAL DESCRIPTION**

This ecosystem consists of open grasslands found in valley bottoms, adjacent to rivers and streams, and on south facing slopes. The ecosystem tends to be mesic to submesic and occurs on loamy fluvial parent material where flooding and/or high water tables increase soil water content and replenish nutrients. The soils on these sites tend to have thick Ah horizons.




**SUCCESSIONAL RELATIONSHIPS**

Due to the nature of the site grasslands often remain the climax vegetation on these sites. In the moister lower slope positions shrubs often dominate the site with succession to aspen and spruce. Disturbance regime, cold air drainage, and competition from a diverse cover of shrubs, forbs and grasses slow or inhibit the establishment of trees. If trees do become established, the rich loamy soils usually result in rapid growth.

**INDICATOR SPECIES**

- Rough fescue
- California oatgrass
- Tufted hairgrass
- Sedge
- Bearberry
- Strawberry
- Three flowered avens
- Clover
- Alpine rough fescue
- Shrubby cinquefoil
- Slender wheatgrass
- Kentucky bluegrass
- Hairy wildrye



**SITE CHARACTERISTICS**

- Moisture regime: mesic, submesic
- Nutrient regime: rich, medium
- Topographic position: crest, upper, mid to lower slope
- Slope: (0-5%) (5-20%)
- Aspect: south, southwest


**SOIL CHARACTERISTICS**

- Organic thickness: (0-5)
- Humus form: moder, mull
- Surface texture: CL, SiL, L
- Effective texture: CL, SiL, SL
- Depth to Mottles/Gley: none
- Drainage: well, moderately well, imperfectly
- Parent material: F, C, E, GF
- Soil subgroup: O.EB ,O.HR, CU.R

**ECOSITE PHASES**

- ff1 grassland (44)
- ff2 shrubland (26)

**ff1 Grassland (n=44)**



**CHARACTERISTIC SPECIES**

**Shrub**

- [ 3 ] Bearberry
- [ 4 ] Shrubby cinquefoil

**Forb**

- [ 1 ] Slender blue beard tongue
- [ 1 ] Graceful cinquefoil
- [ 9 ] Three flowered avens
- [ 2 ] Wild strawberry

**Grasses**

- [ 24 ] Rough fescue
- [ 2 ] Tufted hairgrass
- [ 4 ] Hairy wildrye
- [ 9 ] California oatgrass
- [ 19 ] Sedge
- [ 6 ] Slender wheatgrass
- [ 2 ] Alpine rough fescue

**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
 Nutrient regime: rich  
 Topographic position: midslope, lowerslope, level  
 Slope: 5-20%  
 Aspect: southerly


**SOIL CHARACTERISTICS**

Organic thickness: (0-5)  
 Humus form: mull, moder  
 Surface texture: CL, SiL, L  
 Effective texture: CL, SiL, SL  
 Depth to Mottles/Gley: none  
 Drainage: well, modetately well, imperfectly  
 Parent material: E, C, F, GF  
 Soil subgroup: O.HR, O.EB, CU.R

**RANGE PLANT COMMUNITY TYPES**

- UFA5. Rough fescue-Tufted hairgrass n=5
- UFA6. Rough fescue-Hairy wildrye n=11
- UFA7. Rough fescue-California oatgrass/Bearberry n=4
- UFA7a. California oatgrass-Rough fescue/Bearberry n=2
- UFA8. California oatgrass-Sedge n=10
- UFA12. Rough fescue-Bog sedge n=9
- UFA13. Arctic rough fescue n=2
- UFA16. Hairy wildrye-Rough fescue/Bearberry n=1

**ff2 Shrubland (n=26)**



**SITE CHARACTERISTICS**

Moisture regime: mesic, subhygric,hygric  
 Nutrient regime: rich  
 Topographic position: lowerslope, level  
 Slope: 0-10%  
 Aspect: south westerly, north easterly

**SOIL CHARACTERISTICS**

Organic thickness: (0-4)  
 Humus form: moder  
 Surface texture: L  
 Effective texture: CL  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: L, M  
 Soil subgroup: O.EB

**RANGE PLANT COMMUNITY TYPES**

**CHARACTERISTIC SPECIES**

**Shrub**

- [ 6 ] Willow
- [ 17 ] Bog birch
- [ 4 ] Bearberry

**Forb**


- [ 1 ] Alpine bistort
- [ 5 ] Strawberry
- [ 1 ] Lindley's aster
- [ 2 ] Fireweed

**Grasses**

- [ 1 ] Hairy wildrye
- [ 1 ] Sheep fescue
- [ 4 ] Sedge
- [ 6 ] California oatgrass
- [ 12 ] Rough fescue
- [ 1 ] Purple oatgrass
- [ 3 ] Slender wheatgrass

- UFB4. Willow/Rough fescue n=2
- UFB5. Bog birch/Rough fescue/Bearberry n=18
- UFB6. Willow/California oatgrass-Sedge n=4
- UFB8. Willow/Hairy wildrye-Sedge n=2

**ff1a Grazed grassland (n=57)**



**CHARACTERISTIC SPECIES**

**Shrub**

- [ 1 ] Shubby cinquefoil
- [ 1 ] Bog birch

**Forb**

- [ 2 ] Strawberry
- [ 2 ] Yarrow
- [ 1 ] Graceful cinquefoil
- [ 19 ] Clover

**Grasses**

- [ 5 ] Sedge
- [ 2 ] Idaho fescue
- [ 2 ] Parry's oatgrass
- [ 32 ] Creeping red fescue
- [ 1 ] Rocky mountain fescue
- [ 15 ] Kentucky bluegrass

**SITE CHARACTERISTICS**

Moisture regime: mesic, submesic  
 Nutrient regime: rich  
 Topographic position: midslope, lowerslope, level  
 Slope: 5-20%  
 Aspect: southerly


**SOIL CHARACTERISTICS**

Organic thickness: (0-5)  
 Humus form: mull, moder  
 Surface texture: CL, SiL, L  
 Effective texture: CL, SiL, SL  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well, imperfectly  
 Parent material: E, C, F, GF  
 Soil subgroup: O.HR, O.EB, CU.R

**RANGE PLANT COMMUNITY TYPES**

- UFC2. Rocky Mtn. fescue/Graceful cinquefoil n=1
- UFC7. Creeping red fescue/Clover n=24
- UFC10. Purple oatgrass-Rough fescue n=1
- UFC11. Slender wheatgrass-Sedge-Rough fescue(n=29)
- UFA17. Idaho fescue-Parry oatgrass-Sedge n=2

**ff2a Grazed shrubland (n=2)**



**SITE CHARACTERISTICS**

Moisture regime: mesic, subhygric,hygric  
 Nutrient regime: rich  
 Topographic position: lowerslope, level  
 Slope: 0-10%  
 Aspect: south westerly, north easterly

**SOIL CHARACTERISTICS**

Organic thickness: (0-4)  
 Humus form: moder  
 Surface texture: L  
 Effective texture: CL  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: L, M  
 Soil subgroup: O.EB

**RANGE PLANT COMMUNITY TYPES**

UFC9. Willow/Kentucky bluegrass n=2

**CHARACTERISTIC SPECIES**

**Shrub**

[22] Willow


**Forb**

- [11] Dandelion
- [7] Yarrow
- [2] Strawberry
- [5] Tall lungwort
- [5] Clover

**Grasses**

- [12] Kentucky bluegrass
- [5] Sheep fescue
- [9] Slender wheatgrass
- [8] Tufted hairgrass

**f4b Harvested bracted honeysuckle/  
Sw (n=1)**



**SITE CHARACTERISTICS**

**Moisture regime:** subhygric, mesic  
**Nutrient regime:** rich  
**Topographic position:** midslope, upper slope, level  
**Slope:** (20)%  
**Aspect:** southeasterly

**SOIL CHARACTERISTICS**

**Organic thickness:** (6-15), (16-25)  
**Humus form:** raw moder  
**Surface texture:** Si, L, SiL, SiCL, SiC  
**Effective texture:** C, SiCL, SiC, Si, L, CL  
**Depth to Mottles/Gley:** none  
**Drainage:** Moderately well, imperfectly  
**Parent material:** M, L, E, C/M  
**Soil subgroup:** D.GL, O.GL, E.EB, E.DYB

**RANGE PLANT COMMUNITY TYPES**

UFF5. River alder-Willow/Fireweed-Cow parsnip n=1

**CHARACTERISTIC SPECIES**

**Tree**

- [ 3 ] White spruce
- [ 5 ] Aspen

**Shrub**

- [ 5 ] Willow
- [ 5 ] River alder

**Forb**

- [ 21 ] Fireweed
- [ 13 ] Cow parsnip
- [ 10 ] Stinging nettle
- [ 10 ] White geranium
- [ 8 ] Tall lungwort
- [ 6 ] Horsetail


**Grasses**

- [ 4 ] Slender wheatgrass
- [ 3 ] Marsh reedgrass

**g2a Grazed forb meadow (n=4)**

**SITE CHARACTERISTICS**

Moisture regime: submesic, subhygric  
 Nutrient regime: medium, rich  
 Topographic position: midslope, level  
 Slope: (0-6)%  
 Aspect: southeast, southwest



**SOIL CHARACTERISTICS**

Organic thickness: (6-15)  
 Humus form: mull  
 Surface texture: SiC, SiL, C  
 Effective texture: SiC, C  
 Depth to Mottles/Gley: none  
 Drainage: moderately well, imperfectly  
 Parent material: L, F  
 Soil subgroup: R.G, O.R

**RANGE PLANT COMMUNITY TYPES**

UFC8. K. Bluegrass-Timothy/Meadow rue(n=4)

**CHARACTERISTIC SPECIES**

**Shrub**

[ 2 ] Willow

**Forb**

[ 13 ] Dandelion

[ 7 ] Cow parsnip

[ 5 ] Clover

[ 7 ] Veiny meadow rue


**Grasses**

[ 7 ] Slender wheatgrass

[ 2 ] Smooth brome

[ 17 ] Timothy

[ 22 ] Kentucky bluegrass



Moisture regime: mesic, subhygric,hygric  
 Nutrient regime: very rich  
 Topographic position: lowerslope, level  
 Slope: 0%, 5-40%  
 Aspect: south westerly, south easterly, south and east

**SOIL CHARACTERISTICS**

Organic thickness: (0-5)  
 Humus form: moder  
 Surface texture: SiL, L  
 Effective texture: SiL, LS, L  
 Depth to Mottles/Gley: none  
 Drainage: well, moderately well  
 Parent material: GL, GF, F  
 Soil subgroup: O.DYB, O.HR

**RANGE PLANT COMMUNITY TYPES**

UFA2. Sedge-Slender wheatgrass/Meadow rue n=3  
 UFA3. Tufted hairgrass-Sedge n=35  
 UFA4. Tufted hairgrass-Sedge-Slender wheatgrass n=8

**CHARACTERISTIC SPECIES**

**Shrub**

- [ 1 ] Barclay's willow
- [ 2 ] Willow
- [ 2 ] Bog birch


**Forb**

- [ 7 ] Lindley's aster
- [ 8 ] Yarrow
- [ 8 ] Veiny meadow rue
- [ 5 ] Graceful cinquefoil
- [ 6 ] Wild strawberry

**Grasses**

- [ 36 ] Sedge species
- [ 3 ] Slender wheatgrass
- [ 25 ] Tufted hairgrass

**g3a Grazed grass meadow (n=74)**



**CHARACTERISTIC SPECIES**

**Shrub**

[ 1 ] Willow

**Forb**

- [ 4 ] Strawberry
- [ 8 ] Yarrow
- [ 9 ] Graceful cinquefoil
- [ 15 ] Dandelion
- [ 5 ] Clover
- [ 7 ] Veiny meadow rue

**Grasses**

- [ 21 ] Sedge species\*
- [ 7 ] Slender wheatgrass
- [ 8 ] Tufted hairgrass
- [ 28 ] Kentucky bluegrass
- [ 1 ] Fringed brome

**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric,hygric

**Nutrient regime:** very rich

**Topographic position:** lowerslope, level

**Slope:** 0%, 5-40%

**Aspect:** south westerly, south easterly, south and east

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-5)

**Humus form:** moder

**Surface texture:** SiL, L

**Effective texture:** SiL, LS, L

**Depth to Mottles/Gley:** none

**Drainage:** moderately well, poor

**Parent material:** GL, GF, F

**Soil subgroup:** O.DYB, O.HR

**RANGE PLANT COMMUNITY TYPES**

UFC1. Slender wheatgrass-Sedge/Low forbs n=9


UFC3. Kentucky bluegrass/Dandelion n=23

UFC4. Kentucky bluegrass-Sedge/Dandelion n=25

UFC5. Tufted hairgrass-Kentucky bluegrass n=4

UFC6. Sedge/Tufted hairgrass n=13

## j2 Horsetail Pb (n=1)



## SITE CHARACTERISTICS

Moisture regime: hygric, subhygric, subhydric  
 Nutrient regime: rich, very rich, medium, poor  
 Topographic position: toe, midslope, lowerslope, level  
 Slope: level, (2-%)  
 Aspect: level, northerly, westerly

## SOIL CHARACTERISTICS

Organic thickness: (6-15)(0-5)  
 Humus form: raw moder, mor, peatymor  
 Surface texture: SiL, L, SL  
 Effective texture: SiL, LS, L  
 Depth to Mottles/Gley: none, (26-50)(0-25)  
 Drainage: imperfect, poor, moderately well  
 Parent material: F, C  
 Soil subgroup: R.G, O.R, O.EB, GL.EB

## RANGE PLANT COMMUNITY TYPES

UFD6. Pb/Willow/Horsetail (n=1)

## CHARACTERISTIC SPECIES

### Trees

[ 35 ] Balsam poplar  
 [ 5 ] Aspen  
 [ 3 ] White spruce

### Shrub

[ 30 ] Willow  
 [ 3 ] Rose


### Forb

[ 12 ] Horsetail  
 [ 9 ] Scouring rush  
 [ 3 ] Tall lungwort  
 [ 1 ] Scouring rush  
 [ 4 ] Clover  
 [ 7 ] Strawberry

### Grasses

[ 1 ] Hairy wildrye  
 [ 1 ] Marsh reedgrass  
 [ 1 ] Kentucky bluegrass

**j1b Harvested horsetail/ Sw (n=1)**



**CHARACTERISTIC SPECIES**

**Forb**

- [ 18 ] Dandelion
- [ 3 ] Strawberry
- [ 22 ] Yarrow
- [ 8 ] Graceful cinquefoil
- [ 5 ] Veiny meadow rue

**Grasses**

- [ 4 ] Creeping red fescue
- [ 4 ] Slender wheatgrass
- [ 46 ] Kentucky bluegrass

**SITE CHARACTERISTICS**

Moisture regime: hygric, subhygric  
 Nutrient regime: rich  
 Topographic position: level, midslope, lower slope, toe  
 Slope: (1)%  
 Aspect: north

**SOIL CHARACTERISTICS**

Organic thickness: (6-15), (0-5)  
 Humus form: mor, raw moder, peatymor  
 Surface texture: SiL, SL, L  
 Effective texture: SiL, LS, L  
 Depth to Mottles/Gley: none  
 Drainage: imperfectly, moderately well, poor, very poor  
 Parent material: F, C  
 Soil subgroup: R.G, O.R, O.EB, GL.EB

**RANGE PLANT COMMUNITY TYPES**

UFF3. Sw/Horsetail/Kentucky bluegrass(n=1)

**UPPER FOOTHILLS SUBREGION**  
**NATIVE GRASSLANDS AND SHRUBLANDS**



**Figure 1. Overview of native shrub and grassland complex in the Upper Foothills subregion**



## Native grass and shrublands

The native grass and shrubland community types (Table 2) are found in the valley bottoms, adjacent to streams and rivers, throughout the Upper Foothills subregion. Deep snow accumulations and cold air drainage prevent trees from growing in these valley bottoms (Daubenmire, 1978). Historically, these grass and shrublands burned frequently, further preventing tree encroachment.

The sequence of these community types along a moisture gradient from wet (UFA1 sedge meadows) to dry (UFA9 junegrass-sedge/ sage slopes) is outlined in Figure 3. The change in species composition from the wet sedge meadows to rough fescue and California oatgrass meadows may occur over a 3 foot elevational gradient.

The maintenance of these grassland community types is extremely fire dependent. The lack of fire allows bog birch and willow to expand, shading the modal grassland community types. Prolonged shading causes the understory composition to shift from a tufted hairgrass-rough fescue dominated understory to one dominated by slender wheatgrass and sedge (Figure 3). Under heavy shrub cover (pussy willow shrubland and willow-bog birch community types), there is little forb or grass understory. Increased shrub cover also causes a decline in forage productivity and reduces the accessibility for livestock.



**Figure 2. Typical native shrub and grassland transition zone in Upper Foothills subregion**



**Table 2.** Forage production and ecosite phases of the native grass and shrubland community types for the Upper Foothills subregion  
 Community number      Community type      Grass      Forb      Productivity (kg/ha)      Moisture      Drainage      Carrying capacity (Ha/AUM)

Community number	Community type	Grass	Forb	Productivity (kg/ha)	Moisture	Drainage	Carrying capacity (Ha/AUM)
<b>a1</b>	<b>Ecosite phase shrubby grassland</b>	<b>400</b>			<b>Subseric</b>	<b>Rapidly</b>	<b>Non-use</b>
UFA9.	Junegrass-Sedge/ Sage	400		824	Subseric	Rapidly	Non-use
UFA10.	Bearberry/ Slender wheatgrass	400		171	Subseric	Well	Non-use
<b>c6</b>	<b>Ecosite phase hairy wildrye grassland</b>	<b>1115</b>	<b>312</b>	<b>8</b>	<b>Submesic</b>	<b>Mod. Well</b>	<b>2.1</b>
UFA15.	Hairy wildrye-Sedge	222	66	8	Submesic	Well	3.1
UFA16.	Hairy wildrye-Bearberry	2008	557		Mesic	Mod. Well	1.0
<b>ff1</b>	<b>Ecosite phase grassland</b>				<b>Mesic</b>	<b>Well</b>	<b>0.9</b>
UFA5.	Rough fescue-Tufted hairgrass	1068	618		Subhygric	Mod. Well	0.5
UFA6.	Rough fescue-Hairy wildrye	2041	358		Mesic	Well	0.4
UFA7.	Rough fescue-Bearberry	1023	538		Submesic	Well	0.6
UFA7a	California oatgrass-Rough fescue/ Bearberry	1023	538		Submesic	Well	Non-use
UFA8.	California oatgrass-Sedge	1051	373	585	Mesic	Well	0.6
UFA12.	Rough fescue-Bog sedge	966	149		Mesic	Well	0.8
UFA13.	Arctic Rough fescue	743	372		Subhygric	Mod. Well	0.8
UFA17	Idaho fescue-Parry's oatgrass-Sedge	1053	372	44	Mesic	Well	3.1
<b>ff2</b>	<b>Ecosite phase shrubland</b>	<b>1057</b>	<b>235</b>	<b>446</b>	<b>Subhygric</b>	<b>Mod. Well</b>	<b>0.6</b>
UFB4.	Willow/ Rough fescue	640	132	733	Subhygric	Mod. Well	0.6
UFB5.	Bog birch/ Rough fescue/ Bearberry	1206	173	306	Subhygric	Well	0.6
UFB6.	Willow/ California oatgrass-Sedge	1324	400	300	Subhygric	Mod. well	0.5
UFB8.	Willow/ Hairy wildrye-Sedge				Subhygric	Mod. well	0.5
<b>f6</b>	<b>Ecosite phase bracted honeysuckle willow</b>	<b>162</b>	<b>1786</b>		<b>Hygric</b>	<b>Imperfectly</b>	<b>Non-use</b>
UFB12	Alder-Willow/ Horsetail	162	1786		Hygric	Imperfectly	Non-use

<b>g1 Ecosite phase shrubby meadow</b>													
UFB2.	Willow/ Slender wheatgrass-Sedge	1148	437	566	1487	Subhygric	Mod. Well	Non-use					
UFB3.	Willow/ Tufted hairgrass	1573	735		1669	Subhygric	Well	0.5					
UFB7.	Pussy willow shrubland	1168	405	772	2058	Subhygric	Mod. Well	0.4					
UFB9.	Bog birch/ Sedge-Marsh reedgrass	796	58	322	181	Subhygric	Mod. well	Non-use					
UFB10.	Willow-Bog birch/ Sedge	937	176	730	1176	Hygric	Imperfectly	Non-use					
UFB11.	Willow-Bog birch	1265	811	438	1730	Hygric	Imperfectly	Non-use					
<b>g2 Ecosite phase forb meadow</b>		200	1154	400	3126	Subhygric	Mod. well	0.7					
UFA11.	Fireweed/ Hairy wildrye				1252	Subhygric	Mod. Well	0.7					
UFA14.	Cow parsnip-Meadow rue/ Fringed brome				5000	Subhygric	Mod. Well	0.7					
<b>g3 Ecosite phase grass meadow</b>		1748	742	277	2495	Subhygric	Mod. Well	0.4					
UFA2.	Sedge-Slender wheatgrass/ Meadow rue				2500*	Subhygric	Mod. Well	0.4					
UFA3.	Tufted hairgrass-Sedge	1665	513	277	2241	Subhygric	Mod. Well	0.4					
UFA4.	Tufted hairgrass-Sedge-Slender wheatgrass	1831	971		2745	Sughygric	Mod. Well	0.3					
<b>m2 Ecosite phase shrubby rich fen</b>		1325	126	732	2105	Subhydric	Poorly	Non-use					
UFB1.	Willow-Bog birch/ Water sedge	1325	126	732	2105	Subhydric	Poorly	Non-use					
<b>m3 Ecosite phase graminoid rich fen</b>		1981	384	872	2381	Subhydric	Poorly	Non-use					
UFA1.	Water sedge meadows	1981	384	872	2381	Subhydric	Poorly	Non-use					

\* Estimated

Arctic rough fescue  
 Rough fescue-Bog sedge  
 (higher elevations)




Figure 3. Layout of plant community types for native grass and shrublands in the landscape of the Upper Foothills subregion.

**Key to Grassland Community Types (Grasslands dominated by native species)**

1. Hydric, periodic flooding, depressional, dominated by sedges ..... UFA1 Water sedge meadows  
 Drier, ungrazed or lightly grazed dominated by forbs or grass (rough fescue, tufted hairgrass) ..... 2
2. Moist sites dominated by Forbs (Fireweed, Cow parsnip) ..... 3  
 Grass dominated (upland sedges, rough fescue, slender wheatgrass, hairy wildrye, tufted hairgrass) ..... 4
3. Moist, lowland sites dominated by fireweed ..... UFA11 Fireweed/ Hairy wildrye (Forb meadow)  
 Fine textured, silty soils, dominated by cow parsnip ..... UFA14 Cow parsnip-Meadow rue/ Fringed brome
4. Moist sites dominated by Tufted hairgrass with no rough fescue present at site, includes moderately grazed sites  
 dominated by slender wheatgrass, Rocky Mtn. fescue and sedge species ..... 5  
 Drier sites dominated by rough fescue, hairy wildrye, california oatgrass, slender wheatgrass ..... 6
5. Early successional sites with little slender wheatgrass, dominated by Tufted hairgrass  
 ..... UFA3 Tufted hairgrass-Sedge  
 Later successional or grazed tufted hairgrass or rough fescue meadows with abundant forbs, sedge, and  
 slender wheatgrass ..... 5a  
 5a Lightly grazed sites dominated by tufted hairgrass, Sedge and Slender wheatgrass....  
 .....UFA4 Tufted hairgrass-Sedge-Slender wheatgrass  
 Lightly to moderately grazed sites dominated by sedge, slender wheatgrass, Rocky Mtn. fescue or  
 graceful cinquefoil.....5b  
 5b Moderately to heavily grazed site dominated by Rocky Mtn. fescue or graceful cinquefoil.....  
 .....UFC2 Rocky Mtn. fescue/Graceful cinquefoil  
 Lightly to moderately grazed site dominated by slender wheatgrass, sedge, low forb species or rough fescue  
 .....5c  
 5c Dry well drained sites with some rough fescue present.....UFC1 Sedge-Slender wheatgrass-Rough fescue  
 Moister sites, tufted hairgrass present.....UFC1 Slender wheatgrass-Sedge/Low forbs
6. Rough fescue dominated, higher elevation and moist sites ..... 7  
 Hairy wildrye, California oatgrass, slender wheatgrass, Idaho fescue, Parry oatgrass or Purple oatgrass  
 dominated ..... 12
7. Moist Alpine sites with globeflower, fleabane, monkshood, mountain heliotrope or drier sites with bog sedge .. 8  
 Drier sites at lower elevations tufted hairgrass, hairy wildrye, bearberry or slender wheatgrass codominant .... 9
8. Dry well drained sites with bog sedge codominant ..... UFA12 Rough fescue-Bog sedge  
 Moist Alpine sites with alpine forb species ..... UFA13 Alpine rough fescue
9. Moist sites codominated with Tufted hairgrass ..... UFA5 Rough fescue-Tufted hairgrass  
 Drier well drained sites dominated by hairy wildrye, bearberry, ..... 10
10. Lower, south facing slopes, well-developed soils hairy wildrye codominant .. UFA6 Rough fescue-Hairy wildrye  
 Well drained sites with Bearberry present in understory ..... 11
11. Shallow, well drained, gravelly soils, low nutrient ..... UFA7 Rough fescue/ Bearberry  
 Ghost area California oatgrass dominated ..... UFA7a California oatgrass-Rough fescue/ Bearberry
12. California oatgrass dominated, well-drained soil, cold air drainage level areas in valley bottoms  
 ..... UFA8 California oatgrass-Sedge  
 Idaho fescue, Parry oatgrass, Hairy wildrye or Purple oatgrass dominated communities on south facing slopes or  
 dry gravelly river beds ..... 13
13. Low land moist meadows or dry gravelly river beds ..... 15  
 Steep south facing slopes.....14
14. Steep, south facing slopes, shallow soils, drought tolerant species Junegrass, sage UFA9 Junegrass-Sedge/ Sage  
 Hairy wildrye or Idaho fescue, Purple oatgrass, Parry oatgrass dominated sites ..... 16
15. River bed, dry, gravelly, well drained sites ..... UFA10 Bearberry/ Slender wheatgrass  
 Moist meadows dominated by drier sedge species ..... UFA2 Sedge-Slender wheatgrass/ Meadow rue
16. Hairy wildrye dominated ..... 17  
 Moister, south facing slopes in the Ghost area, Idaho fescue, Parry oatgrass, Purple oatgrass dominated  
 grasslands ..... 16a  
 16a Idaho fescue, Parry oatgrass dominated sites in the Ghost area.....UFA17 Idaho fescue-P.oatgrass-Sedge

- Grazed sites dominated by Purple oatgrass and rough fescue.....UFC9 Purple oatgrass-Rough fescue
- 17. South facing slopes, at higher elevation.....UFA15 Hairy wildrye-Sedge
- Grazed site, dominated by hairy wildrye, rough fescue present . . . UFA16 Hairy wildrye-Rough fescue/Bearberry

**Key to Grazing Modified Grassland and Shrubland Community Types** (dominated by non-native species)

- 1. Native dominated ..... 2
- Non-native dominated, creeping red fescue, Kentucky bluegrass, clover, dandelion ... 7
- 2. Grass dominated (slender wheatgrass, rocky mtn. fescue, sedge, purple oatgrass, tufted hairgrass) ..... 3
- Shrub dominated with K. bluegrass understory . . . . . UFC10 Willow/ Kentucky bluegrass
- 3. Moist grassy meadows with rough fescue, tufted hairgrass still present on site ..... 4
- Drier, well drained sites dominated by Rocky mtn. fescue
- ..... UFC2 Rocky mtn. Fescue/ Graceful cinquefoil
- 4. Slender wheatgrass, Sedge and purple oatgrass dominated community.....4a
- Grazed tufted hairgrass communities(moister sites).....5
- 4a Slender wheatgrass and Sedge dominated.....UFC1 Slender wheatgrass-Sedge/Low forbs
- Purple oatgrass dominates.....UFC9 Purple oatgrass-Rough fescue
- 3. Kentucky bluegrass present . . . . . UFC5 Tufted hairgrass-Kentucky bluegrass
- Kentucky bluegrass absent, recovering site . . . . . UFC6 Sedge-Tufted hairgrass
- 4. Kentucky bluegrass dominated ..... 8
- Seeded sites with Creeping red fescue . . . . . UFC7 Creeping red fescue/ Clover
- 5. Heavily grazed sites with dandelion as co-dominant ..... UFC3 Kentucky bluegrass/ Dandelion
- ..... UFC3 Kentucky bluegrass/ Dandelion
- Heavily grazed cow parsnip meadow, lower elevation sites, cow parsnip present in small amounts . . . . . UFC8 Kentucky bluegrass-Timothy/ Meadow rue

**Key to Shrubland Community Types**

1. Bog birch dominated ..... 2  
 Willow dominated ..... 3

2. Dry, nutrient poor soils, with rough fescue ..... UFB5 Bog birch/ Rough fescue/ Bearberry  
 Wet, poor drainage, no willow cover ..... UFB9 Bog birch/ Sedge-Marsh reedgrass

3. Poorly drained sites with water sedge in understory, or shrublands with little understory ... 4  
 Drier well drained sites with slender wheatgrass, hairy wildrye, rough fescue, tufted hairgrass  
 found in the understory ..... 7

4. Poor drainage, very wet sites ..... UFB1 Willow-Bog birch/ Water sedge  
 Shrub dominated sites with little understory willow and bog birch dominated ..... 5

5: Tall willow or alder dominated ..... 6  
 Short willow, imperfectly drained sites little understory ..... UFB11 Willow-Bog birch

6. Occurring along water bodies ..... UFB7 Pussy willow shrubland  
 Moist, nutrient rich seepage sites with alder and willow .... UFB12 Alder-Willow/ Horsetail

7. Well drained sites with Hairy wildrye dominating the understory, typical of well drained valley  
 bottomland sites ..... UFB8 Willow/ Hairy wildrye-Sedge  
 Recently invaded grasslands with rough fescue, tufted hairgrass or california oatgrass,  
 slender wheatgrass or graceful sedge ..... 8

8. California oatgrass dominated ..... UFB6 Willow/ California oatgrass-Sedge  
 Rough fescue or tufted hairgrass present in understory ..... 9

9. Rough fescue dominated ..... UFB4 Willow/ Rough fescue  
 Tufted hairgrass, graceful sedge or slender wheatgrass dominated ..... 10

10. Tufted hairgrass dominated ..... UFB3 Willow/ Tufted hairgrass  
 Sedge or Slender wheatgrass dominated understory.....11

11. Sedge dominated ..... UFB10 Willow-Bog birch/ Sedge  
 Slender wheatgrass dominated ..... UFB2 Willow/ Slender wheatgrass-Sedge

**UFA1. Water-Beaked sedge meadows**  
*(Carex aquatilis-Carex rostrata)*

n=14 Wet conditions and periodic flooding result in the formation of water sedge meadows. Bog birch and willow will invade into the drier edges of these meadows to form the willow-bog birch/ water sedge community type (UFB1).

These community types are quite productive, producing nearly 2000 kg/ha of forage, but the high water table in the spring and summer when these meadows are most palatable limits livestock use. A study in the Yukon found that crude protein on these meadows declined from a high of 10% in May to less than 5% in September (Bailey et al., 1992). As a result, these meadows would be rated as secondary or non-use range.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp</i> )	2	0-10	79
<b>BOG BIRCH</b>			
( <i>Betula glandulosa</i> )	1	0-1	29
<b>FORBS</b>			
<b>ARROW LEAVED COLTSFOOT</b>			
( <i>Petasites sagittatus</i> )	2	0-20	7
<b>GRASSES</b>			
<b>WATER SEDGE</b>			
( <i>Carex aquatilis</i> )	14	0-63	29
<b>BEAKED SEDGE</b>			
( <i>Carex rostrata</i> )	2	0-30	7
<b>SEDE</b>			
( <i>Carex spp</i> )	64	0-96	50
<b>TUFTED HAIRGRASS</b>			
( <i>Deschampsia cespitosa</i> )	11	0-40	86

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYDRIC  
 NUTRIENT REGIME  
 MESOTROPHIC  
 ELEVATION:  
 1091-1760 M (1484 M)  
 SOIL DRAINAGE:  
 POORLY  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1981 (810-4438)  
 FORBS: 384 (46-776)  
 SHRUBS: 872 (8-1736)  
 TOTAL: 2381 (966-4684)

SUGGESTED GRAZING CAPACITY  
 NON-USE

## UFA2. Sedge-Slender wheatgrass/ Meadow rue (*Carex spp.-Agropyron trachycaulum/ Thalictrum venulosum*)

n=3 These sites are dominated by three sedge species that are adapted to moist conditions: *C. praticola*, *C. praegracilis* and *C. prairea*. The presence of small amounts of tufted hairgrass and rough fescue indicates that these sites may represent a phase of the rough fescue-tufted hairgrass plant community (UFA5). Past heavy grazing pressure may have shifted the plant community to one dominated by sedge species or these sites could be too wet for tufted hairgrass and rough fescue growth.

The forage productivity on this community type is good. In comparison with the water sedge meadows (UFA1), these sites remain drier throughout the growing season and this allows easy access for livestock. This community would be rated as primary range.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	0-2	67
BOG BIRCH			
( <i>Betula glandulosa</i> )	1	0-2	33
<b>FORBS</b>			
VEINY MEADOW RUE			
( <i>Thalictrum venulosum</i> )	28	20-36	100
OLD MAN'S WHISKERS			
( <i>Geum triflorum</i> )	9	0-14	67
SLENDER BLUE BEARDTONGUE			
( <i>Penstemon procerus</i> )	5	0-8	67
YARROW			
( <i>Achillea millefolium</i> )	10	5-14	100
SILVERY CINQUEFOIL			
( <i>Potentilla arguta</i> )	5	0-8	67
LINDLEY'S ASTER			
( <i>Aster ciliolatus</i> )	2	0-5	33
<b>GRASSES</b>			
MEADOW SEDGE			
( <i>Carex praticola</i> )	16	0-26	67
GRACEFUL SEDGE			
( <i>Carex praegracilis</i> )	11	0-32	33
PRAIRIE SEDGE			
( <i>Carex prairea</i> )	9	0-28	33
SEDEGE			
( <i>Carex spp</i> )	25	0-75	33
SLENDER WHEATGRASS			
( <i>Agropyron trachycaulum</i> )	8	0-12	67

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1460 M  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
EAST  
SLOPE:  
0-5%

### FORAGE PRODUCTION(KG/HA)

TOTAL: 2500

<p><b>SUGGESTED GRAZING CAPACITY</b> 0.4 HA/AUM OR 0.8 ACRES/AUM</p>
--

**UFA3. Tufted hairgrass-Sedge**  
(*Deschampsia cespitosa-Carex praegracilis*)

n=34 This community is located on moist sites that are better drained and slightly drier than the pure sedge meadows (UFA1). Willoughby (1992) and Willoughby (1999), found that tufted hairgrass is a common plant species on these lowland sites throughout the Upper Foothills and Lower Subalpine subregions. At lower elevations, this species appears to be replaced by marsh reedgrass. When this community type is protected from grazing for 25-30 years, willow and bog birch expand (willow/ tufted hairgrass (UFB3)) and tufted hairgrass and sedge decline (Willoughby, 1992). The decline in graminoid cover also results in a decline in available forage production from 2200 to 1800 kg/ha. Continuous heavy grazing pressure causes hairgrass to decline and the site will be invaded by Kentucky bluegrass and dandelion.

Bork (1994), found this community type. to be the most productive type described in Willmore wilderness park. Forage production averages over 2000 kg/ha and can vary from 800-3300 kg/ha. This community type would be rated as primary range.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

**SHRUBS**

BARCLAY'S WILLOW ( <i>Salix barclayi</i> )	1	0-8	23
WILLOW ( <i>Salix spp</i> )	1	0-12	14
BOG BIRCH ( <i>Betula glandulosa</i> )	1	0-10	23
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	8	0-41	94
STRAWBERRY ( <i>Fragaria virginiana</i> )	7	0-27	77
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	6	0-23	74
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	7	0-44	57
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	6	0-23	89
DANDELION ( <i>Taraxacum officinale</i> )	5	0-30	60
<b>GRASSES</b>			
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	31	2-70	100
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	12	0-43	69
WATER SEDGE ( <i>Carex aquatilis</i> )	1	0-20	11
SEDGE ( <i>Carex spp</i> )	26	0-88	14

SLENDER WHEATGRASS

(*Agropyron trachyaulum*) 9 0-27 71

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1276-1800 M (1461 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
VARIABLE  
SLOPE:  
0-40%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1665 (422-3654)  
FORBS: 513 (6-1572)  
SHRUBS: 277 (124-414)  
TOTAL: 2455 (824-3706)

<p><b>SUGGESTED GRAZING CAPACITY</b> 0.4 HA/AUM OR 0.9 ACRES/AUM</p>
--

**UFA4. Tufted hairgrass-Sedge-Slender wheatgrass**  
*(Deschampsia cespitosa-Carex spp.-Agropyron trachycaulum)*

n=8 This community type may be transitional between the willow dominated community types and the tufted hairgrass dominated grasslands. Two of the sites described in this community are represented by the inside, ungrazed transect at two rangeland reference area sites. Protection from grazing for 25-35 years appears to allow willow to expand and there is a shift away from a tufted hairgrass dominated community type to a type that is dominated by slender wheatgrass, sedge and tall forb species. Continued protection from grazing and fire will likely lead to a community dominated by willow and bog birch with little understory of forbs and grass.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
BARCLAY'S WILLOW <i>(Salix barclayi)</i>	3	0-13	38
WILLOW <i>(Salix spp)</i>	6	0-33	38
BOG BIRCH <i>(Betula glandulosa)</i>	5	0-26	50
<b>FORBS</b>			
LINDLEY'S ASTER <i>(Aster ciliolatus)</i>	6	0-15	50
VEINY MEADOW RUE <i>(Thalictrum venulosum)</i>	11	0-31	88
YARROW <i>(Achillea millefolium)</i>	7	T-13	100
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	4	T-13	100
STRAWBERRY <i>(Fragaria virginiana)</i>	6	0-15	88
AMERICAN VETCH <i>(Vicia americana)</i>	3	1-5	100
FIREWEED <i>(Epilobium angustifolium)</i>	4	0-8	75
TALL LUNGWORT <i>(Mertensia paniculata)</i>	8	0-32	88
<b>GRASSES</b>			
GRACEFUL SEDGE <i>(Carex praegracilis)</i>	11	0-21	75
TUFTED HAIRGRASS <i>(Deschampsia cespitosa)</i>	9	1-17	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	13	0-28	88

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1303-1805 M (1423 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1831 (864-2416)  
 FORBS: 971 (477-1702)  
 TOTAL: 2745 (1478-4118)

<p>SUGGESTED GRAZING CAPACITY                  0.3 HA/AUM OR 0.7 ACRES/AUM</p>
--

## UFA5. Rough fescue-Tufted hairgrass

(*Festuca scabrella-Deschampsia cespitosa*)

n=5 This community type is located up slope from the tufted hairgrass-sedge community type (UFA3) on drier, better drained soils. The drier soil conditions limit the amount of forage being produced. There was 300 kg/ha less forage produced in the rough fescue-tufted hairgrass community type compared to the tufted hairgrass-sedge community type (UFA3).

In the absence of fire and grazing, this community type will become dominated by willow and bog birch (willow/ rough fescue community type (UFB4)). Heavy grazing pressure also decreases the cover of rough fescue and tufted hairgrass and allows Kentucky bluegrass and dandelion to increase (Willoughby, 1992). The dominant plant species on this community are highly palatable and the sites are easily accessible to livestock. Consequently, this community would be rated as primary range.

### PLANT COMPOSITION CANOPY COVER(%)

	Mean	Range	Const.
<b>FORBS</b>			
SLENDER BLUE BEARDTONGUE ( <i>Penstemon procerus</i> )	4	1-9	100
YARROW ( <i>Achillea millefolium</i> )	4	1-11	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	3	0-6	80
CHICKWEED ( <i>Cerastium arvense</i> )	1	0-4	80
MONKSHOOD ( <i>Aconitum delphinifolium</i> )	1	0-4	40
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	5	0-25	40
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca scabrella</i> )	23	18-28	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	17	3-29	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	4	1-6	100
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	13	0-50	60
CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	3	0-8	60

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1370-1737 M (1532 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
SOUTHEAST  
SLOPE:  
0-3%

### FORAGE PRODUCTION (KG/HA)

GRASS: 1068 (605-1797)  
FORBS: 618 (166-1252)  
TOTAL: 1684 (913-2272)

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM OR 1.2 ACRES/AUM

**UFA6. Rough fescue-Hairy wildrye**  
(*Festuca scabrella*-*Elymus innovatus*)

n=18 These grasslands are located on lower, south facing slopes. They represent the transition zone from the dry junegrass-sedge/ sedge (UFA9) dominated south facing slopes to the moist rough fescue and tufted hairgrass dominated community types (UFA5). Grazing pressure causes a shift away from a rough fescue, hairy wildrye dominated community (UFA6) to a sedge, Kentucky bluegrass dominated community (UFC4) (Willoughby, 1992). These grasslands are fairly moist and have well developed soils which makes them very productive. This community type would be rated as primary range.

This community type is very similar to the rough fescue dominated communities described in the Ya Ha Tinda, west of Sundre (Willoughby et al. 2001)

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
BEBB'S WILLOW ( <i>Salix bebbiana</i> )	2	0-13	18
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	0-13	78
<b>FORBS</b>			
FIREWEED ( <i>Epilobium angustifolium</i> )	3	0-30	27
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	5	0-19	56
STAR FLOWERED SOLOMON'S SEAL ( <i>Smilacina stellata</i> )	2	0-27	33
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	4	0-9	67
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	4	0-19	73
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca scabrella</i> )	35	8-85	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	8	0-28	67
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	4	0-18	61
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	1	0-5	39
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	5	0-42	22
SEDGE ( <i>Carex spp</i> )	9	0-24	44
PRAIRIE SEDGE ( <i>Carex prairea</i> )	1	0-18	11
JUNEGRASS ( <i>Koeleria macrantha</i> )	4	0-19	67

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1320-1798 M (1589 M)  
SOIL DRAINAGE:  
WELL  
ASPECT:  
SOUTH  
SLOPE:  
5-48%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 2041 (472-4832)  
FORBS: 358 (12-696)  
TOTAL: 2190 (484-5162)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM OR 0.9 ACRES/AUM

**UFA7. Rough fescue/ Bearberry**  
*(Festuca scabrella/ Arctostaphylos uva-ursi)*

n=4 This community type was described in the Upper Clearwater Forest Land Use Zone and is similar to the bog birch/ rough fescue/ bearberry community type but lacks the cover of bog birch. Willoughby (2001) felt that bog birch indicated sites with deeper snow accumulations. This community occupies sites that have shallow, well-drained, and gravelly soils and there maybe little snow cover which does not favour the growth of bog birch. This community is moderately productive but because of the poor soil conditions, precautions must be taken to prevent overutilization

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
<b>BEARBERRY</b>			
<i>(Arctostaphylos uva-ursi)</i>	21	12-31	100
<b>SHRUBBY CINQUEFOIL</b>			
<i>(Potentilla fruticosa)</i>	2	0-5	75
<b>FORBS</b>			
<b>OLD MAN'S WHISKERS</b>			
<i>(Geum triflorum)</i>	10	0-26	75
<b>YARROW</b>			
<i>(Achillea millefolium)</i>	11	0-38	75
<b>GRACEFUL CINQUEFOIL</b>			
<i>(Potentilla gracilis)</i>	3	0-10	50
<b>GRASSES</b>			
<b>ROUGH FESCUE</b>			
<i>(Festuca scabrella)</i>	49	38-56	100
<b>SLENDER WHEATGRASS</b>			
<i>(Agropyron trachycaulum)</i> 9		5-19	100
<b>GRACEFUL SEDGE</b>			
<i>(Carex praegracilis)</i>	3	0-6	75
<b>FRINGED BROME</b>			
<i>(Bromus ciliatus)</i>	4	0-7	75
<b>HAIRY WILD RYE</b>			
<i>(Elymus innovatus)</i>	4	0-9	75

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBXERIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1676-1829 M (1745 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTH  
 SLOPE:  
 0-5%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 1023 (580-1686)  
 FORBS: 538 (204-820)  
 TOTAL: 1561 (1156-1890)

**SUGGESTED GRAZING CAPACITY**  
 0.6 HA/AUM OR 1.3 ACRES/AUM

**UFA7a. California oatgrass-Rough fescue/ Bearberry**  
*(Danthonia californica-Festuca scabrella/ Arctostaphylos uva-ursi)*

n=2 This community type was described in the Ghost area west of Calgary. It appears to represent a transitional grassland between the Montane and Upper Foothills subregions. This community also appears to be transitional between the California oatgrass dominated grasslands (UFA8) and the previously described Rough fescue/Bearberry dominated community. This community is located on steep, south facing slopes and small hillcrests with well-drained subxeric soils. The dry site conditions limit the amount of forage available for domestic livestock and the steep slopes restrict livestock access. Consequently, this community type should be rated non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	21	4-36	100
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	8	1-15	100
<b>FORBS</b>			
OLD MAN'S WHISKERS <i>(Geum triflorum)</i>	20	11-27	100
YARROW <i>(Achillea millefolium)</i>	2	T-3	100
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	1	T-1	100
<b>GRASSES</b>			
CALIFORNIA OATGRASS <i>(Danthonia californica)</i>	29	27-30	100
ROUGH FESCUE <i>(Festuca scabrella)</i>	15	7-22	100
IDAHO FESCUE <i>(Festuca idahoensis)</i>	8	5-11	100
SEDGE <i>(Carex spp)</i>	11	4-7	100
HAIRY WILDRYE <i>(Elymus innovatus)</i>	1	0-2	50

MOISTURE REGIME:

SUBXERIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1745 M

SOIL DRAINAGE:

WELL

ASPECT:

SOUTH

SLOPE:

25%

**FORAGE PRODUCTION (KG/HA)**

TOTAL 1561 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
0.6 HA/AUM OR 1.3 AC/AUM

**ENVIRONMENTAL VARIABLES**

**UFA8. California oatgrass-Sedge**  
(*Danthonia californica*-*Carex praegracilis*)

n=9 Dry, gravelly or stony soils support this moderately productive grassland that is dominated by California oatgrass. Small pockets of this community type occur throughout the Upper Foothills subregion. In the Yukon, these small meadows were found to form in depressions which appeared to act as pronounced frost pockets (Bailey et al., 1992). In the Subalpine subregion, these California oatgrass dominated grasslands are often associated with bog sedge (Willoughby 1999). The cold air drainage and poor nutrient quality of the soil limits the forage productivity of these sites.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	7	0-25	60
DWARF BILBERRY ( <i>Vaccinium caespitosum</i> )	1	0-5	10
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	2	0-10	30
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	8	2-15	100
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	14	0-46	90
ALPINE MILK VETCH ( <i>Astragalus alpinus</i> )	3	0-17	30
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	7	0-25	90
BLUE EYED GRASS ( <i>Sisyrinchium montanum</i> )	2	0-19	30
<b>GRASSES</b>			
CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	31	0-57	90
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	10	0-37	50
SEEDGE ( <i>Carex spp</i> )	18	0-50	50
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	8	0-36	80
SHEEP FESCUE ( <i>Festuca saximontana</i> )	3	0-15	40
COLUMBIA NEEDLEGRASS ( <i>Stipa columbiana</i> )	4	0-21	30

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1400-1580 M (1484 M)

SOIL DRAINAGE:

WELL

ASPECT:

VARIABLE

SLOPE:

0-45%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 1051 (400-1582)

FORBS: 373 (118-762)

SHRUBS: 585 (110-1402)

TOTAL: 1578 (1033-2014)

**SUGGESTED GRAZING CAPACITY**  
0.6 HA/AUM OR 1.3 ACRES/AUM

**ENVIRONMENTAL VARIABLES**

**UFA9. Junegrass-Sedge/ Sage**  
*(Koeleria macrantha-Carex spp./ Artemisia frigida)*

n=4 This community type occurs on steep, south facing slopes with shallow soils overlying sandstone bedrock. The majority of the vegetation is composed of the drought tolerant species: sage, bearberry and junegrass. The inaccessibility and fragile nature of the soils make this community type unsuitable for grazing.

This community type is very similar to the blunt sedge-rocky mtn. fescue/ bearberry community described by Willoughby (1999) and the junegrass-hairy wildrye-brome community described by Corns and Achuff (1982) on steep south-facing slopes in the Subalpine subregion.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
PRICKLY ROSE <i>(Rosa acicularis)</i>	2	0-3	75
<b>BEARBERRY</b> <i>(Arctostaphylos uva-ursi)</i>	1	0-5	25
<b>FORBS</b>			
PLAINS WORMWOOD <i>(Artemisia campestris)</i>	1	0-5	25
FRINGED SAGE <i>(Artemisia frigida)</i>	7	0-17	75
MOUNTAIN GOLDENROD <i>(Solidago spathulata)</i>	1	0-5	25
LATE YELLOW LOCOWEED <i>(Oxytropis monticola)</i>	1	0-3	25
<b>GRASSES</b>			
JUNEGRASS <i>(Koeleria macrantha)</i>	19	13-30	100
THREAD-LEAVED SEDGE <i>(Carex filifolia)</i>	14	0-32	50
SEDEGE <i>(Carex spp)</i>	14	0-38	75
SHEEP FESCUE <i>(Festuca saximontana)</i>	1	0-5	25

**MOISTURE REGIME:**

SUB XERIC

**NUTRIENT REGIME:**

SUBMESOTROPHIC

**ELEVATION:**

1560-1720 M (1592 M)

**SOIL DRAINAGE:**

RAPIDLY

**ASPECT:**

SOUTH

**SLOPE:**

25-40%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 737 (400-1044)

FORBS: 359 (222-495)

SHRUBS: 171 (1-400)

TOTAL: 1147 (800-1378)

SUGGESTED GRAZING CAPACITY

NON-USE

**ENVIRONMENTAL VARIABLES**

**UFA10. Bearberry/ Slender wheatgrass**  
*(Arctostaphylos uva-ursi/ Agropyron trachycaulum)*

n=2 This community type is found scattered throughout the Upper Foothills subregion on dry, gravelly, well drained river flats. The presence of silverberry, yellow mountain avens, bearberry and early yellow locoweed are very common on these sites.

The poor soil conditions limit the forage productivity and amount of regrowth after grazing. This community type should be rated as secondary or non-use range.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
<b>SILVERBERRY</b>			
<i>(Elaeagnus commutata)</i>	1	0-1	50
<b>YELLOW MOUNTAIN AVENS</b>			
<i>(Dryas drummondii)</i>	2	0-3	50
<b>BEARBERRY</b>			
<i>(Arctostaphylos uva-ursi)</i>	11	8-12	100
<b>FORBS</b>			
<b>STRAWBERRY</b>			
<i>(Fragaria virginiana)</i>	22	14-29	100
<b>EARLY YELLOW LOCOWEED</b>			
<i>(Oxytropis sericea)</i>	12	0-24	50
<b>YARROW</b>			
<i>(Achillea millefolium)</i>	3	1-5	100
<b>GRASSES</b>			
<b>JUNEGRASS</b>			
<i>(Koeleria macrantha)</i>	3	0-5	50
<b>ROUGH FESCUE</b>			
<i>(Festuca scabrella)</i>	4	0-8	50
<b>SLENDER WHEATGRASS</b>			
<i>(Agropyron trachycaulum)</i>	6	0-11	100
<b>ALPINE BLUEGRASS</b>			
<i>(Poa alpina)</i>	5	0-10	50
<b>SHEEP FESCUE</b>			
<i>(Festuca saximontana)</i>	2	0-3	50

MOISTURE REGIME:  
 SUBMESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1400-1415 M (1408 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTHEAST  
 SLOPE:  
 0-3%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 400  
 SHRUBS: 400  
 TOTAL: 500

SUGGESTED GRAZING CAPACITY  
 NON-USE

**ENVIRONMENTAL VARIABLES**

## UFA11. Fireweed/ Hairy wildrye (Forb meadow)

(*Epilobium angustifolium*/ *Elymus innovatus*)

n=3 This community type is found on moist lowland sites adjacent to the lodgepole pine and white spruce dominated forests. It represents the transition from the willow and grass dominated riparian areas to the conifer dominated forests. In the absence of disturbance (fire) it appears that succession of conifers into the grassy meadows shifts the species dominance away from a predominantly graminoid cover to one dominated by forbs such as fireweed, Lindley's aster and palmate leaved coltsfoot. There is also a shift in grass cover away from tufted hairgrass, rough fescue and sedge species to more shade tolerant grass species such as purple oatgrass and hairy wildrye. Periodic burning of this site is required to limit tree and shrub expansion.

This community type is very productive and easily accessible to livestock. It would be rated as primary range.

### PLANT COMPOSITION CANOPY COVER(%)

	Mean	Range	Const.
<b>UNDERSTORY TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> )	3	0-8	67
LODGEPOLE PINE ( <i>Pinus contorta</i> )	6	0-10	67
<b>SHRUBS</b>			
WILLOW ( <i>Salix spp.</i> )	16	0-25	67
<b>FORBS</b>			
FIREWEED ( <i>Epilobium angustifolium</i> )	25	T-47	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	7	3-13	100
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	16	0-26	67
YARROW ( <i>Achillea millefolium</i> )	7	3-11	100
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	10	T-20	100
PURPLE OATGRASS ( <i>Schizachne purpurascens</i> )	6	0-15	67
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	3	0-4	67

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1310-1454 M (1401 M)

SOIL DRAINAGE:

MODERATELY WELL

ASPECT:

SOUTHEAST

SLOPE:

0-2%

### FORAGE PRODUCTION (KG/HA)

GRASS: 200

FORBS: 1154

SHRUBS: 400

TOTAL: 1252 (1182-1354)

SUGGESTED GRAZING CAPACITY

0.7 HA/AUM OR 1.6 AC/AUM

### ENVIRONMENTAL VARIABLES

## UFA12. Rough fescue-Bog sedge

*(Festuca scabrella-Kobresia myosuroides)*

n=2 This community is very similar to the bog birch/ rough fescue-bog sedge community type described by Willoughby (1999) in the Foothills ecodistrict of the Subalpine subregion. Bog sedge is well adapted to growing on dry alpine slopes and rocky ridges in the mountains. Corns and Achuff (1982), described bog sedge dominated community types on windswept ridges in the alpine subregion of Banff and Jasper National Parks. The sites described in this community type appear to represent the transition from the Upper Foothills to the Subalpine subregion.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL

*(Potentilla fruticosa)* 1 0-1 50

BEARBERRY

*(Arctostaphylos uva-ursi)* 6 0-11 50

FORBS

FIREWEED

*(Epilobium angustifolium)* 1 0-1 50

OLD MAN'S WHISKERS

*(Geum triflorum)* 19 15-21 100

FALSE DANDELION

*(Agoseris glauca)* 3 1-6 100

ALPINE HEDYSARUM

*(Hedysarum alpinum)* 1 0-5 22

GRASSES

HAIRY WILDRYE

*(Elymus innovatus)* 10 2-17 100

ROUGH FESCUE

*(Festuca scabrella)* 38 32-43 100

BOG SEDGE

*(Kobresia myosuroides)* 25 12-37 100

SLENDER WHEATGRASS

*(Agropyron trachycaulum)* 18 12-22 100

SEDGE

*(Carex spp.)* 8 2-13 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1432-1828 M (1688 M)

SOIL DRAINAGE:

WELL

ASPECT:

SOUTHERLY

SLOPE:

0-40%

### FORAGE PRODUCTION (KG/HA)

GRASS: 966 (832-1234)

FORBS: 149 (98-202)

TOTAL: 1115 (932-1434)

SUGGESTED GRAZING CAPACITY

0.8HA/AUM

## UFA13. Arctic rough fescue

(*Festuca altaica*)

n=2 This community was described at higher elevations in Willmore Wilderness Park. Bork (1994), described this community type on alpine and subalpine slopes where climate and soil conditions are still suitable for fescue to dominate in the stand. The community has a subhygric moisture regime and is moderately well drained. Forb species such as globeflower, fleabane, monkshood, and mountain heliotrope are all characteristic of these high elevation meadows.

This community is much wetter than the rough fescue-bog sedge community (UFA12) previously described and is similar to the forb meadows community type described by Willoughby (1999) in the Subalpine subregion.

### PLANT COMPOSITION CANOPY COVER(%)

	Mean	Range	Const.
<b>SHRUBS</b>			
DWARF BILBERRY ( <i>Vaccinium caespitosum</i> )	3	2-3	100
DWARF BIRCH ( <i>Betula glandulosa</i> )	1	0-1	50
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	2	1-3	100
MONKSHOOD ( <i>Aconitum delphinifolium</i> )	1	0-2	100
MOUNTAIN HELIOTROPE ( <i>Valeriana sitchensis</i> )	1	0-2	100
WANDERING DAISY ( <i>Erigeron peregrinus</i> )	2	0-3	50
GLOBEFLOWER ( <i>Trollius albiflorus</i> )	2	0-4	50
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca altaica</i> )	47	36-57	100
MOUNTAIN TIMOTHY ( <i>Phleum commutatum</i> )	2	2-2	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	2	2-2	100
SEDGE ( <i>Carex spp.</i> )	6	5-7	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	2	0-4	50

MOISTURE REGIME:

SUBHYRGIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1510-2000 M (1755 M)

SOIL DRAINAGE:

MODERATELY WELL

### FORAGE PRODUCTION (KG/HA)

GRASS: 743 (527-959)

FORBS: 372 (368-375)

TOTAL: 1115 (895-1334)

### SUGGESTED GRAZING CAPACITY

0.8 HA/AUM OR 1.7 AC/AUM

### ENVIRONMENTAL VARIABLES

## UFA14. Cow parsnip-Veiny meadow rue/ Fringed brome

(*Heracleum lanatum*-*Thalictrum venulosum*/ *Bromus ciliatus*)

n=1 This community type is transitional between the Lower Foothills and Upper Foothills subregions. It was described on fine textured, silty soils adjacent to the Baptiste river west of Rocky Mountain House. Grazed stands of this community type were also described in the Solomon valley, west of Hinton. Increased grazing pressure generally allows timothy, Kentucky bluegrass and dandelion to increase with a corresponding drop in the cover of cow parsnip, meadow rue and the native grasses and sedges. The high moisture and nutrient regime of this site makes it extremely productive, and once it has been invaded by agronomic species it is highly palatable for domestic livestock. It is difficult to find representative stands of this community type that have not been grazed.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW	-	-	100
( <i>Salix spp.</i> )	4	-	100
SNOWBERRY			
( <i>Symphoricarpos</i>			
<i>occidentalis</i> )	4	-	100
<b>FORBS</b>			
COW PARSNIP			
( <i>Heracleum lanatum</i> )	21	-	100
VEINY MEADOW RUE			
( <i>Thalictrum venulosum</i> )	10	-	100
TALL LUNGWORT			
( <i>Mertensia paniculata</i> )	11	-	100
TALL LARKSPUR			
( <i>Delphinium glaucum</i> )	8	-	100
FIREWEED			
( <i>Epilobium angustifolium</i> )	5	-	100
AMERICAN VETCH			
( <i>Vicia americana</i> )	3	-	100
<b>GRASSES</b>			
KENTUCKY BLUEGRASS			
( <i>Poa pratensis</i> )	15	-	100
AWNED SEDGE			
( <i>Carex atherodes</i> )	7	-	100
SLENDER WHEATGRASS			
( <i>Agropyron trachycaulum</i> )	4	-	100
SEDEGE			
( <i>Carex spp.</i> )	12	-	100
FRINGED BROME			
( <i>Bromus ciliatus</i> )	6	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1060 M

SOIL DRAINAGE:

MODERATELY WELL

### FORAGE PRODUCTION (KG/HA)

TOTAL: 5000

**SUGGESTED GRAZING CAPACITY**

0.3 HA/AUM OR 0.4 AC/AUM

## UFA15. Hairy wildrye-Sedge

(*Elymus innovatus*-*Carex* spp.)

n=1 This community type was described on south facing slopes along Wilson Creek in the Upper Foothills subregion. Corns and Achuff (1982) described similar community types in the subalpine of Banff and Jasper National Parks. These included the shrubby cinquefoil/ hairy wildrye and hairy wildrye/ bearberry-juniper community types. Both of these community types were associated with steep south facing slopes. The presence of this community type may indicate the transition to the Subalpine subregion.

This community type does not produce a large amount of forage because of the dry site conditions and poor nutrient content of the soil, but the lack of open areas for livestock grazing in this subregion makes these grassland sites attractive to livestock.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### FORBS

SHOWY LOCOWEED			
( <i>Oxytropis splendens</i> )	18	-	100
AMERICAN VETCH			
( <i>Vicia americana</i> )	8	-	100
LOW GOLDENROD			
( <i>Solidago multiradiata</i> )	6	-	100
STRAWBERRY			
( <i>Fragaria virginiana</i> )	2	-	100
FIREWEED			
( <i>Epilobium angustifolium</i> ) <sup>4</sup>		-	100
ALPINE HEDYSARUM			
( <i>Hedysarum alpinum</i> )	1	-	100
GRASSES			
HAIRY WILDRYE			
( <i>Elymus innovatus</i> )	50	-	100
ARCTIC BLUEGRASS			
( <i>Poa arctica</i> )	6	-	100
SEDGE			
( <i>Carex</i> spp.)	5	-	100

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

SUBMESIC

#### NUTRIENT REGIME:

SUBMESOTROPHIC

#### ELEVATION:

1860 M

#### SOIL DRAINAGE:

WELL

#### ASPECT:

SOUTHWEST

#### SLOPE:

5%

### FORAGE PRODUCTION(KG/HA)

GRASS: 222

FORBS: 66

SHRUBS: 8

TOTAL: 296

### SUGGESTED GRAZING CAPACITY

3.1 HA/AUM OR 6.9 AC/AUM

## UFA16. Hairy wildrye-Rough fescue/ Bearberry

*(Elymus innovatus-Festuca scabrella/ Arctostaphylos uva-ursi)*

n=1 This community was described in the Ghost area west of Calgary on a well drained, level valley floor. It appears to represent a grazed rough fescue, bearberry or hairy wildrye dominated community. Willoughby (2000) found that heavy grazing on the rough fescue dominated grasslands often leads to a community that is dominated by sedge and hairy wildrye. Protection from grazing or a reduction in stocking rate allows this community type to recover back to a rough fescue dominated community. The time frame for complete recovery takes 20 years (Willoughby 2000)

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

<b>SHRUBS</b>			
BEARBERRY			
<i>(Arctostaphylos uva-ursi)</i>	7	-	100
SHRUBBY CINQUEFOIL			
<i>(Potentilla fruticosa)</i>	1	-	100
<b>FORBS</b>			
FIREWEED			
<i>(Epilobium angustifolium)</i>	5	-	100
GRACEFUL CINQUEFOIL			
<i>(Potentilla gracilis)</i>	4	-	100
GOLDENROD			
<i>(Solidago missouriensis)</i>	1	-	100
SMOOTH ASTER			
<i>(Aster laevis)</i>	1	-	100
STRAWBERRY			
<i>(Fragaria virginiana)</i>	13	-	100
VEINY MEADOW RUE			
<i>(Thalictrum venulosum)</i>	1	-	100
<b>GRASSES</b>			
SLENDER WHEATGRASS			
<i>(Agropyron trachycaulum)</i>	2	-	100
HAIRY WILDRYE			
<i>(Elymus innovatus)</i>	11	-	100
SEDGE			
<i>(Carex spp.)</i>	1	-	100
ROUGH FESCUE			
<i>(Festuca scabrella)</i>	2	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1680

SOIL DRAINAGE:

MODERATELY WELL

ASPECT:

EAST

SLOPE:

3%

### FORAGE PRODUCTION(KG/HA)

GRASS: 2008

FORBS: 557

TOTAL: 2121

**SUGGESTED GRAZING CAPACITY**

0.4 HA/AUM OR 1.0 AC/AUM

## UFA17. Idaho fescue-Parry oatgrass-Sedge

(*Festuca idahoensis*-*Danthonia parryi*-*Carex spp.*)

n=2 This community type was described in the Ghost area west of Calgary. This area represents a transition between the Montane and Upper Foothills subregions. This community type is very similar to moderately and heavily grazed rough fescue dominated communities in the Montane subregion. Both Idaho fescue and Parry oatgrass are more characteristic of the Montane subregion and increase with increased grazing pressure. Protection from grazing will often allow this community type to recover back to a Rough fescue-Parry oatgrass dominated community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	2-3	100
BOG BIRCH ( <i>Betula glandulosa</i> )	3	0-5	50
WILLOW ( <i>Salix spp</i> )	2	0-4	50
<b>FORBS</b>			
STRAWBERRY ( <i>Fragaria virginiana</i> )	22	10-33	100
YARROW ( <i>Achillea millefolium</i> )	13	6-18	100
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	10	7-11	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	4	0-8	50
<b>GRASSES</b>			
SEDGE ( <i>Carex spp.</i> )	39	28-50	100
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	22	28-50	100
PARRY'S OATGRASS ( <i>Danthonia parryi</i> )	21	16-25	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	2	0-3	50

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1400 M

SOIL DRAINAGE:

IMPERFECTLY

ASPECT:

SOUTH

SLOPE:

2-15%

### FORAGE PRODUCTION(KG/HA)

TOTAL: 1467 \*ESTIMATE

SUGGESTED GRAZING CAPACITY

HA/AUM OR AC/AUM

0.6 HA/AUM OR 1.4 AC/AUM

**UFB1. Willow-Bog birch/ Water sedge**  
*(Salix spp.-Betula glandulosa/ Carex aquatilis)*

n=45 This shrub community appears in areas with very poor drainage. It is found in association with the wetter water sedge meadows (UFA1). These sites are fairly productive but are difficult to graze due to the moist ground conditions and heavy shrub cover which reduces access and mobility in the area. Increased flooding and prolonged water logging may result in the disappearance of willow and a transition to a water sedge meadow.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
BARCLAY'S WILLOW <i>(Salix barclayi)</i>	2	0-48	9
WILLOW <i>(Salix spp.)</i>	28	0-65	84
BOG BIRCH <i>(Betula glandulosa)</i>	11	0-58	56
<b>FORBS</b>			
ARROW LEAVED COLTSFOOT <i>(Petasites sagittatus)</i>	1	0-13	36
LINDLEY'S ASTER <i>(Aster ciliolatus)</i>	1	0-6	11
STICKY PURPLE GERANIUM <i>(Geranium viscosissimum)</i>	1	0-14	11
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	1	0-7	29
ARCTIC ASTER <i>(Aster sibiricus)</i>	1	0-8	2
<b>GRASSES</b>			
WATER SEDGE <i>(Carex aquatilis)</i>	11	0-76	24
SEEDGE <i>(Carex spp.)</i>	40	0-82	71
TUFTED HAIRGRASS <i>(Deschampsia cespitosa)</i>	6	0-35	80
MARSH REEDGRASS <i>(Calamagrostis canadensis)</i>	1	0-13	11

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :  
 SUBHYDRIC  
 NUTRIENT REGIME  
 MESOTROPHIC  
 ELEVATION:  
 1227-1820 M (1443 M)  
 SOIL DRAINAGE:  
 POORLY  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-14%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 1325 (340-3000)  
 FORBS: 126 (2-402)  
 SHRUBS: 732 (54-2180)  
 TOTAL: 2105 (814-4662)

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFB2. Willow/ Slender wheatgrass-Sedge**  
*(Salix spp./ Agropyron trachycaulum-Carex spp.)*

n=4 This community type is very similar to the tufted hairgrass-sedge-slender wheatgrass c.t. (UFA4) previously described. Both community types appear to represent the various stages of succession onto tufted hairgrass meadows. When these communities are protected from disturbance (fire and grazing), willow and bog birch expand and tufted hairgrass declines. Willow growth also appears to favour the growth of tall forbs (veiny meadow rue, fireweed, aster) and slender wheatgrass. Fire has played a dominant role in controlling brush encroachment in the past and continued protection will allow continued shrub expansion, resulting in a decline in forage production.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

BARCLAY'S WILLOW  
*(Salix barclayi)* 13 0-19 75

BOG BIRCH  
*(Betula glandulosa)* 13 2-23 100

**FORBS**

VEINY MEADOW RUE  
*(Thalictrum venulosum)* 7 3-9 100

OLD MAN'S WHISKERS  
*(Geum triflorum)* 17 3-23 100

STRAWBERRY  
*(Fragaria virginiana)* 13 10-15 100

TALL LARKSPUR  
*(Delphinium glaucum)* 1 0-T 50

LINDLEY'S ASTER  
*(Aster ciliolatus)* 10 0-26 75

YARROW  
*(Achillea millefolium)* 8 4-13 100

**GRASSES**

GRACEFUL SEDGE  
*(Carex praegracilis)* 24 0-37 75

SEDEGE  
*(Carex spp.)* 7 0-29 25

SLENDER WHEATGRASS  
*(Agropyron trachycaulum)* 15 0-22 75

CALIFORNIA OATGRASS  
*(Danthonia californica)* 7 0-20 100

TUFTED HAIRGRASS  
*(Deschampsia cespitosa)* 4 0-12 75

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1349-1615 M (1455 M)

SOIL DRAINAGE:

WELL

ASPECT:

NORTH TO NORTHEAST

SLOPE:

0-4%

**FORAGE PRODUCTION (KG/HA)**

GRASS: 1573

FORBS: 735

TOTAL: 1669 (900-2308)

<p><b>SUGGESTED GRAZING CAPACITY</b>          0.5 HA/AUM OR 1.2 ACRES/AUM</p>
---

**UFB3. Willow/ Tufted hairgrass**  
(*Salix barclayi/ Deschampsia cespitosa*)

n=16 This community type is found in association with the tufted hairgrass-sedge c.t. (UFA3). Willow encroachment into a tufted hairgrass meadow eventually results in this community type. Historically, fire has played an important role in the maintenance of the grassland community types in this subregion. Continued fire suppression will eventually allow willow and bog birch to invade many of these grassy meadows.

The encroachment of willow onto the tufted hairgrass-sedge c.t. causes a decline in forage production from 2200 kg/ha to 1200 kg/ha for grass and forb production. Continued protection of this community type from disturbance will most likely lead to the development of a community type similar to the willow/ slender wheatgrass (UFB2) and then to the pussy willow shrubland (UFB7). The latter community has a high cover of willow (71%) and very little forage for domestic livestock.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>SHRUBS</b>			
BARCLAY'S WILLOW ( <i>Salix barclayi</i> )	33	0-85	88
BOG BIRCH ( <i>Betula glandulosa</i> )	11	0-75	88
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	6	2-14	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	9	0-30	94
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	11	0-25	81
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	5	0-21	81
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	4	0-10	88
DANDELION ( <i>Taraxacum officinale</i> )	2	0-11	50
<b>GRASSES</b>			
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	18	0-38	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	8	0-25	94
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	12	0-31	69
PURPLE OATGRASS ( <i>Schizachne purpurascens</i> )	4	0-32	56

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1104-1667 M (1434 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
VARIABLE  
SLOPE:  
0-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1168 (275-2484)  
FORBS: 405 (8-1052)  
SHRUBS: 772 (36-2250)  
TOTAL: 2058 (500-5132)

**SUGGESTED GRAZING CAPACITY**  
0.4 HA/AUM OR 1.0 ACRES/AUM

**UFB4. Willow/ Rough fescue**  
(*Salix barclayi/ Festuca scabrella*)

n=2 This community type was described by Bork (1994) in Willmore Wilderness Park. Bork felt this community type originated from recent shrub encroachment onto rough fescue grasslands. Continued shrub expansion will result in decreasing forage productivity. Bork also felt that fescue will be replaced by wheatgrass and sedge plant species. These plants being better adapted to shading and competition from adjacent shrubs.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**TREES**

SUBALPINE FIR -  
(*Abies lasiocarpa*)      1      0-1      50

**SHRUBS**

WILLOW  
(*SALIX* spp)      26      25-27      100

**BOG BIRCH**

(*Betula glandulosa*)      24      10-38      100

**YELLOW MOUNTAIN AVENS**

(*Dryas drummondii*)      3      0-5      50

**ALPINE BEARBERRY**

(*Arctostaphylos rubra*)      4      0-7      50

**FORBS**

ALPINE BISTORT  
(*Polygonum viviparum*)      10      T-19      100

**MONKSHOOD**

(*Aconitium delphinifolium*)      3      0-5      50

**INDIAN PAINTBRUSH**

(*Castilleja miniata*)      2      0-4      50

**GRASSES**

ROUGH FESCUE  
(*Festuca scabrella*)      13      12-14      100

**GRACEFUL SEDGE**

(*Carex praegracilis*)      2      0-3      50

**ENVIRONMENTAL VARIABLES**

**MOISTURE REGIME:**

HYGRIC

**NUTRIENT REGIME:**

PERMESOTROPHIC

**ELEVATION:**

1465-1560 M (1528 M)

**SOIL DRAINAGE:**

MODERATELY WELL

**ASPECT:**

WESTERLY

**SLOPE:**

0-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 640 (600-680)

FORBS: 132 (64-200)

SHRUBS: 733 (150-1316)

TOTAL: 1505 (950-2060)

**SUGGESTED GRAZING CAPACITY**

0.6 HA/AUM OR 1.3 ACRES/AUM

**UFB5. Bog birch/ Rough fescue/ Bearberry**  
*(Betula glandulosa/ Festuca scabrella/ Arctostaphylos uva-ursi)*

n=19 This community type is very similar to the rough fescue-bearberry (UFA7) type previously described, but it is successionaly more advanced. The soils on this community type are gravelly, drier and have a poorer nutrient regime than the other rough fescue and tufted hairgrass dominated community types. The lack of fire on this community type has allowed the shrub cover to expand, reducing forage productivity for wildlife and domestic livestock. In one study, burning a bog birch/ rough fescue/ bearberry community type twice in 3 year intervals controlled birch growth and increased total forage production by over 40% compared to the unburned control (Bork, 1990).

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
BOG BIRCH			
<i>(Betula glandulosa)</i>	33	1-60	100
WILLOW			
<i>(Salix spp)</i>	4	0-13	50
BEARBERRY			
<i>(Arctostaphylos uva-ursi)</i>	9	0-28	90
<b>FORBS</b>			
LINDLEY'S ASTER			
<i>(Aster ciliolatus)</i>	1	0-4	17
ALPINE MILKVETCH			
<i>(Astragalus alpinus)</i>	1	0-9	28
WILD STRAWBERRY			
<i>(Fragaria virginiana)</i>	7	0-24	83
FIREWEED			
<i>(Epilobium angustifolium)</i> 3		0-6	78
SLENDER BLUE BEARDTONGUE			
<i>(Penstemon procerus)</i>	1	0-8	39
OLD MAN'S WHISKERS			
<i>(Geum triflorum)</i>	4	0-17	89
<b>GRASSES</b>			
ROUGH FESCUE			
<i>(Festuca scabrella)</i>	27	3-81	100
GRACEFUL SEDGE			
<i>(Carex praegracilis)</i>	3	0-16	33
SLENDER WHEATGRASS			
<i>(Agropyron trachycaulum)</i> 3		0-20	22
CALIFORNIA OATGRASS			
<i>(Danthonia californica)</i>	8	0-44	72

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1303-1798 M (1562 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTHERLY  
 SLOPE:  
 1-25%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1206 (856-1728)  
 FORBS: 173 (14-394)  
 SHRUBS: 306 (140-582)  
 TOTAL: 1523 (1000-1986)

SUGGESTED GRAZING CAPACITY  
 0.6 HA/AUM OR 1.3 ACRES/AUM

**UFB6. Willow/ California oatgrass-Sedge**  
*(Salix barclayi/ Danthonia californica-Carex spp.)*

n=4 This community type likely develops from willow encroachment onto an oatgrass dominated meadow. The oatgrass meadows are found on dry, gravelly soils. These meadows may also form in frost pockets. The spread of willow is likely caused by the lack of natural disturbance, such as fire. The cover of willow on this community type is fairly extensive. This will restrict access of domestic livestock. This community type would be rated as secondary range.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW SPP.			
<i>(Salix barclayi)</i>	20	0-60	50
<b>BOG BIRCH</b>			
<i>(Betula glandulosa)</i>	10	0-18	75
<b>FORBS</b>			
<b>YARROW</b>			
<i>(Achillea millefolium)</i>	10	2-24	100
<b>GRACEFUL CINQUEFOIL</b>			
<i>(Potentilla gracilis)</i>	6	0-15	75
<b>WILD STRAWBERRY</b>			
<i>(Fragaria virginiana)</i>	20	9-44	100
<b>SLENDER BLUE BEARDTONGUE</b>			
<i>(Penstemon procerus)</i>	2	0-6	50
<b>ALSIKE CLOVER</b>			
<i>(Trifolium pratense)</i>	4	0-17	25
<b>FIREWEED</b>			
<i>(Epilobium angustifolium)</i> 2		0-4	100
<b>VEINY MEADOW RUE</b>			
<i>(Thalictrum venulosum)</i>	4	0-9	75
<b>GRASSES</b>			
<b>CALIFORNIA OATGRASS</b>			
<i>(Danthonia californica)</i>	33	19-56	100
<b>GRACEFUL SEDGE</b>			
<i>(Carex praegracilis)</i>	10	0-30	50
<b>SHEEP FESCUE</b>			
<i>(Festuca saximontana)</i>	7	0-22	50
<b>SLENDER WHEATGRASS</b>			
<i>(Agropyron trachycaulum)</i> 5		0-10	75

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1400-1500 M (1458 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 WEST  
 SLOPE:  
 0-30%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1324 (598-2050)  
 FORBS: 400 (382-418)  
 SHRUBS: 300  
 TOTAL: 1874 (1316-2432)

**SUGGESTED GRAZING CAPACITY**  
 0.5 HA/AUM OR 1.1 ACRES/AUM

**UFB7. Pussy willow shrubland**  
(*Salix discolor*)

n=2 This community type is common along riparian areas, swamps and fringes of marshes and lakes. It appears to be successionaly more advanced than the other willow dominated community types described in this guide. As the willow cover expands over time it shades the understory vegetation resulting in a loss of forage productivity. This community type produces only 200 kg/ha and is generally inaccessible to domestic livestock. This community type should be rated as non-use.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
MEAN RANGE CONST.

<b>SHRUBS</b>			
PUSSY WILLOW ( <i>Salix discolor</i> )	71	70-71	100
<b>BÖG BIRCH</b>			
( <i>Betula glandulosa</i> )	20	8-30	100
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	6	5-7	100
<b>PALMATE LEAVED COLTSFOOT</b>			
( <i>Petasites palmatus</i> )	3	0-0	100
<b>WILD STRAWBERRY</b>			
( <i>Fragaria virginiana</i> )	4	1-7	100
<b>GRASSES</b>			
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	5	1-9	100
<b>MARSH REEDGRASS</b>			
( <i>Calamagrostis canadensis</i> )	3	0-5	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1318-1325 M (1322 M)  
SOIL DRAINAGE:  
MODERATELY WELL

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 181 (101-261)

<p><b>SUGGESTED GRAZING CAPACITY</b> NON-USE</p>
--

**UFB8. Willow/ Hairy wildrye-Sedge**  
(*Salix barclayi/ Elymus innovatus-Carex spp.*)

n=2 This plant community represents a rough fescue-hairy wildrye community type (UFA6) that has continued to undergo succession in the absence of fire and grazing. Willow cover has increased, shading the growth of grasses (rough fescue) and allowing tall-growing forbs (fireweed, aster, veiny meadow rue) to increase. Continued protection from disturbance will allow succession to shrub and eventually tree species. The understorey vegetation will be increasingly shaded and forage production will continue to decrease.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**TREES**

WHITE SPRUCE.

(*Picea glauca*) 1 0-T 50

**SHRUBS**

WILLOW SPP.

(*Salix barclayi*) 30 19-40 100

BOG BIRCH

(*Betula glandulosa*) 21 5-36 100

**FORBS**

LINDLEY'S ASTER

(*Aster ciliolatus*) 12 10-13 100

VEINY MEADOW RUE

(*Thalictrum venulosum*) 6 2-8 100

FIREWEED

(*Epilobium angustifolium*) 5 2-7 100

STRAWBERRY

(*Fragaria virginiana*) 11 8-13 100

TALL LUNGWORT

(*Mertensia paniculata*) 4 0-6 100

**GRASSES**

GRACEFUL SEDGE

(*Carex praegracilis*) 32 20-44 100

HAIRY WILDRYE

(*Elymus innovatus*) 25 13-37 100

PURPLE OATGRASS

(*Schizachne purpurascens*)22 9-35 100

SLENDER WHEATGRASS

(*Agropyron trachycaulum*)22 16-26 100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1371-1400 M (1386) M

SOIL DRAINAGE:

MODERATELY WELL

ASPECT:

NORTHEAST

SLOPE:

0-3%

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 1550 (900-2200)

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM OR 1.1 ACRES/AUM

**UFB9. Bog birch/ Sedge-Marsh reedgrass**  
*(Betula glandulosa/ Carex spp.-Calamagrostis canadensis)*

n=1 This community type occupies valley drainages on soils that are saturated with water for part of the growing season. This type is very similar to the willow-bog birch/ sedge (UFB1) c.t, but lacks the willow cover. It is not clear why there is no willow cover on this type. It is possible that bog birch is better adapted to growing on poor nutrient soils. The presence of marsh reedgrass may indicate the transition from the Lower Foothills to Upper Foothills subregion. Willoughby (1992), observed that marsh reedgrass was more abundant on these lowland sites at lower elevations.

The thick cover of bog birch and very wet conditions restrict access to domestic livestock. Consequently, this community type would be rated as secondary or non-use range.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

<b>SHRUBS</b>			
BOG BIRCH			
<i>(Betula glandulosa)</i>	39	-	100
<b>FORBS</b>			
DWARF RASPBERRY			
<i>(Rubus arcticus)</i>	1	-	100
<b>GRASSES</b>			
MARSH REEDGRASS			
<i>(Calamagrostis canadensis)</i>	11	-	100
TUFTED HAIRGRASS			
<i>(Deschampsia cespitosa)</i>	2	-	100
BROWNISH SEDGE			
<i>(Carex brunnescens)</i>	11	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 HYGRIC  
 NUTRIENT REGIME:  
 SUBMESOTROPHIC  
 ELEVATION:  
 1513 M  
 SOIL DRAINAGE:  
 IMPERFECTLY  
 ASPECT:  
 WEST  
 SLOPE:  
 3%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 796  
 FORBS: 58  
 SHRUBS: 322  
 TOTAL: 1176

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFB10. Willow-Bog birch/ Sedge**  
(*Salix barclayi*-*Betula glandulosa*/ *Carex spp.*)

n=12 This type is very similar to the willow-bog birch/ water sedge community type (UFB1), but the soils are drier and better drained. The drier soil conditions favour the growth of graceful sedge over water sedge.

This community type has a thick cover of bog birch and willow which restricts livestock access to the forage. This community type would be rated as secondary or non-use range.

**PLANT COMPOSITION** CANOPY COVER(%)  
MEAN RANGE CONST.

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
BOG BIRCH ( <i>Betula glandulosa</i> )	21	0-55	75
WILLOW ( <i>Salix spp</i> )	33	2-46	100
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	4	0-15	58
YARROW ( <i>Achillea millefolium</i> )	4	0-11	92
STRAWBERRY ( <i>Fragaria virginiana</i> )	4	0-14	67
TALL LUNGWORT ( <i>Mertensia paniculata</i> )	2	0-6	58
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	3	0-8	67
<b>GRASSES</b>			
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	21	0-53	67
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	4	0-27	58
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	4	0-25	50
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	3	0-10	67

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
HYGRIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1356-1646 M (1523 M)  
SOIL DRAINAGE:  
IMPERFECTLY  
ASPECT:  
VARIABLE  
SLOPE:  
0-15%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 937 (500-1416)  
FORBS: 176 (10-568)  
SHRUB: 730 (200-1522)  
TOTAL: 1730 (800-2468)

SUGGESTED GRAZING CAPACITY  
NON-USE

**UFB11. Willow-Bog birch**  
*(Salix barclayi-Betula glandulosa)*

n=11 This community type is very similar to the willow-bog birch/ sedge c.t. (UFB1), but is successionaly more advanced. The lack of fire has allowed continued expansion of the shrub cover. This has restricted access to livestock and lowered forage productivity. This community type would be rated as non-use for domestic livestock.

**PLANT COMPOSITION** CANOPY COVER(%)  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW SPP <i>(Salix spp)</i>	37	0-85	100
BOG BIRCH <i>(Betula glandulosa)</i>	17	5-50	100
<b>FORBS</b>			
VEINY MEADOW RUE <i>(Thalictrum venulosum)</i>	2	0-9	73
STRAWBERRY <i>(Fragaria virginiana)</i>	5	0-14	73
FIREWEED <i>(Epilobium angustifolium)</i> 2		0-7	73
LINDLEY'S ASTER <i>(Aster ciliolatus)</i>	4	0-16	55
<b>GRASSES</b>			
BALTIC RUSH <i>(Juncus balticus)</i>	2	0-10	90
GRACEFUL SEDGE <i>(Carex praegracilis)</i>	2	0-5	54
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i> 2		0-5	55
TUFTED HAIRGRASS <i>(Deschampsia cespitosa)</i> 2		0-6	64

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1375-1646 M (1472 M)  
 SOIL DRAINAGE:  
 IMPERFECTLY  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1265 (383 -2966)  
 FORBS: 811 (200-1188)  
 SHRUBS: 438 (200-752)  
 TOTAL: 2105 (783-4292)

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFB12. Willow-Alder/ Horsetail**  
*(Salix bebbiana-Alnus tenuifolia/ Equisetum arvense)*

n=1 This community type was described on the boundary between the Upper and Lower foothills subregions in the Solomon valley northwest of Hinton. It is very similar to the to the willow-alder/ shield fern-wild sarsaparilla community described by Lane et al (2000) in the Saddle Hills north of Grande Prairie. This community tends to occupy moist nutrient rich seepage areas which favour the growth of willow, horsetail and fern species. The high cover of willow and alder limits productivity of forbs and grass. It also limits access to domestic livestock. As a result this community type would be rated as non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

BEBB'S WILLOW  
*(Salix bebbiana)* 85 - 100

RIVER ALDER  
*(Alnus tenuifolia)* 15 - 100

BRACED HONEYSUCKLE  
*(Lonicera involcrata)* 5 - 100

**FORBS**

HORSETAIL  
*(Equisetum arvense)* 14 - 100

STRAWBERRY  
*(Fragaria virginiana)* 1 - 100

TALL LUNGWORT  
*(Mertensia paniculata)* 3 - 100

LINDLEY'S ASTER  
*(Aster ciliolatus)* 1 - 100

COW PARSNIP  
*(Heracleum lanatum)* 2 - 100

**GRASSES**

TALL MANNA GRASS  
*(Glyceria grandis)* 1 - 100

KENTUCKY BLUEGRASS  
*(Poa pratensis)* 1 - 100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

HYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1200 M

SOIL DRAINAGE:

IMPERFECTLY

**FORAGE PRODUCTION(KG/HA)**

GRASS: 162

FORBS: 1786

TOTAL: 1948

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UPPER FOOTHILLS SUBREGION**  
**SHRUBLANDS AND GRASSLANDS**  
**MODIFIED BY GRAZING**



**Figure 4. Heavily grazed Kentucky bluegrass/Dandelion community type**



## Grazing modified community types

The grazing modified community types in the Upper Foothills subregion are outlined in Table 3. There are a few grasslands that exhibit signs of historic heavy grazing. These sites are predominantly covered by Kentucky bluegrass, timothy, dandelion and clover plant species (UFC3. Kentucky bluegrass-sedge/ dandelion and UFC4. Kentucky bluegrass/ dandelion and UFC8. Kentucky bluegrass-Timothy/Veiny meadow rue). Under long-term moderate grazing pressure or heavy grazing over a couple of years, there is a general decline in rough fescue and tufted hairgrass and an increase in sedge and slender wheatgrass (UFC1 Slender wheatgrass-sedge/ strawberry and UFC11.Sedge-Slender wheatgrass-Rough fescue). When these plant communities are protected from grazing, they appear to succeed back to the original communities dominated by rough fescue and tufted hairgrass. However, when Kentucky bluegrass becomes established, the community appears to revert to a rough fescue or tufted hairgrass-Kentucky bluegrass-dominated plant community (UFC5. Tufted hairgrass-Kentucky bluegrass) when protected from grazing.

The climax range condition model suggests that vegetation development will be directional, predictable and revert to the original vegetation when protected from grazing, but once Kentucky bluegrass has established, bluegrass appears to compete with rough fescue and tufted hairgrass for co-dominance. These Kentucky bluegrass communities move toward a different community rather than back to the original vegetation when protected from livestock disturbance.

The Rocky Mtn. fescue/ graceful cinquefoil community (UFC2) appears to be a moderately to heavily grazed California oatgrass community type. The dry, gravelly conditions on this site do not appear to favour the growth of Kentucky bluegrass under heavy grazing conditions.

The Creeping red fescue/ Clover (UFC7) community type represents seeded pastures and pipelines within the Upper Foothills subregion. This community type usually occurs at lower elevations, adjacent to farms and ranches where extensive modification of the native grass and shrublands have taken place.

The Purple oatgrass-Rough fescue (UFC9) community type was described in the Ghost area west of Calgary. It appears to represent a rough fescue, hairy wildrye dominated grassland that has undergone heavy grazing pressure. Willoughby (2000) has described purple oatgrass communities on saline soils in the Dry Mixedwood subregion. It is possible that this community type maybe associated with a saline seepage area which favours the growth of purple oatgrass.

Table 3. Shrublands and grasslands modified by heavy grazing pressure in the Upper foothills subregion

Community number	Community type	Production(kg/ha)		Moisture	Drainage	Carrying capacity (Ha/AUM)		
		Grass	Shrub				Total	
<b>f1</b>	<b>Ecosite phase grazed</b>	1789	302	384	2926	Mesic	Well	1.5
UFC2.	Rocky Mtn. fescue-Graceful cinquefoil			384	917	Subhygric	Well	1.0
UFC7.	Creeping red fescue-Clover	1789	302	384	2009	Mesic	Well	0.5
UFC11.	Sedge-Slender wheatgrass-Rough fescue	802	322	18	1207	Mesic	Well	1.1
<b>f2</b>	<b>Ecosite phase grazed</b>	880	316		1196	Subhygric	Mod.well	0.5
UFC10.	Willow/Kentucky bluegrass	880	316		1196	Subhygric	Mod.well	0.5
<b>g2</b>	<b>Ecosite phase forb meadow grazed</b>	2833	1468		4312	Subhygric	Mod. Well	0.3
UFC8.	Kentucky bluegrass-Timothy/Meadow rue	2833	1468		4312	Subhygric	Mod. Well	0.3
<b>g3</b>	<b>Ecosite phase grass meadow grazed</b>	2103	608	212	2744	Subhygric	Mod. Well	0.4
UFC1.	Slender wheatgrass-Sedge/Low forbs	1492	528		1760	Mesic	Mod. Well	0.5
UFC3.	Kentucky bluegrass/Clover-Dandelion	2206	622	150	2837	Mesic	Mod. Well	0.3
UFC4.	Kentucky bluegrass-Sedge/Dandelion	1915	845	102	2767	Mesic	Mod. Well	0.3
UFC5.	Tufted hairgrass-Kentucky bluegrass	3292	1010		4302	Subhygric	Mod. Well	0.2
UFC6.	Sedge-Tufted hairgrass	1661	339		2000	Subhygric	Mod. Well	0.6
UFC9.	Purple oatgrass-Rough fescue	2052	302	384	2798	Submesic	Well	0.3

**UFC1. Slender wheatgrass-Sedge/ Low forbs**  
*(Agropyron trachycaulum-Carex spp./ Fragaria virginiana)*

n=10 This community type appears to arise from grazing a modal fescue-tufted hairgrass community (UFA5). Moderate to heavy grazing causes fescue and hairgrass, both decrease, to decline in the stand. This community is very common in the valley bottoms in areas that are heavily utilized. While still quite productive, these sites have lost two of the most advantageous species. Only a reduction in grazing pressure will once again allow fescue and tufted hairgrass to become prevalent in the stand.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp</i> )	4	0-25	30
<b>FORBS</b>			
STRAWBERRY			
( <i>Fragaria virginiana</i> )	12	0-25	70
YARROW			
( <i>Achillea millefolium</i> )	8	2-14	100
GRACEFUL CINQUEFOIL			
( <i>Potentilla gracilis</i> )	9	0-31	80
LINDLEY'S ASTER			
( <i>Aster ciliolatus</i> )	5	0-20	60
DANDELION			
( <i>Taraxacum officinale</i> )	3	0-8	50
MEADOW RUE			
( <i>Thalictrum venulosum</i> )	4	0-17	70
<b>GRASSES</b>			
SLENDER WHEATGRASS			
( <i>Agropyron trachycaulum</i> )	26	1-58	100
TUFTED HAIRGRASS			
( <i>Deschampsia cespitosa</i> )	1	0-6	10
GRACEFUL SEDGE			
( <i>Carex praegracilis</i> )	21	0-47	80
KENTUCKY BLUEGRASS			
( <i>Poa pratensis</i> )	1	0-4	40
HAIRY WILD RYE			
( <i>Elymus innovatus</i> )	5	0-15	50
FRINGED BROME			
( <i>Bromus ciliatus</i> )	9	0-56	40

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME  
 MESOTROPHIC  
 ELEVATION:  
 1400-2438 M (1623 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 SOUTHERLY  
 SLOPE:  
 0-27%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1492 (824-2548)  
 FORBS: 528 (96-869)  
 TOTAL: 1760 (920-2900)

SUGGESTED GRAZING CAPACITY  
 0.5 HA/AUM OR 1.2 ACRES/AUM

**UFC2. Rocky Mountain fescue/ Graceful cinquefoil**  
*(Festuca brachyphylla/ Potentilla gracilis)*

n=1 This community type was described on a gravelly, well drained site adjacent to Fall creek. It appears that this site was once a California oatgrass-sedge community type (UFA8), but heavy grazing pressure has shifted the community to one dominated by unpalatable low growing graminoids and forbs (Rocky mountain fescue, sedge, yarrow, graceful cinquefoil, pussy toes). The dry site conditions and poor nutrient conditions do not favour the growth of Kentucky bluegrass. This community type would benefit from a deferred rotational grazing system, where the community is rested every other year.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**FORBS**

WILD STRAWBERRY <i>(Fragaria virginiana)</i>	2	-	100
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	13	-	100
YARROW <i>(Achillea millefolium)</i>	8	-	100
ALPINE MILKVETCH <i>(Astragalus alpinus)</i>	6	-	100
ROSEY PUSSY TOES <i>(Antennaria rosea)</i>	2	-	100
RED SEEDED DANDELION <i>(Taraxacum laevigatum)</i>	2	-	100
<b>GRASSES</b>			
ROCKY MOUNTAIN FESCUE <i>(Festuca brachyphylla)</i>	21	-	100
BROWNISH SEDGE <i>(Carex brunnescens)</i>	5	-	100
SLENDER WHEATGRASS <i>(Agropyron trachyalum)</i>	4	-	100
CALIFORNIA OATGRASS <i>(Danthonia californica)</i>	4	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1350 M

SOIL DRAINAGE :

WELL

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 917

**SUGGESTED GRAZING CAPACITY**  
 1.0 HA/AUM OR 2.2 ACRES/AUM

### UFC3. Kentucky bluegrass/ Clover-Dandelion

(*Poa pratensis*/ *Trifolium spp.*-*Taraxacum officinale*)

n=23 This community type develops when the modal tufted hairgrass-sedge dominated communities (UFA3, UFA4) are grazed heavily for prolonged periods of time. Willoughby (1992), felt these grasslands exhibited signs of historic heavy grazing pressure. He felt that under long-term moderate grazing or heavy grazing over a couple of years, rough fescue and tufted hairgrass decline and sedge, slender wheatgrass, and low growing forbs increase. When these plant communities are protected from grazing, they appear to succeed back to the original communities dominated by rough fescue and tufted hairgrass. However, when Kentucky bluegrass becomes established the community appears to revert to a rough fescue or tufted hairgrass-Kentucky bluegrass dominated plant community (UFC5).

These community types are highly productive for domestic livestock during the growing season, but the poor quality of Kentucky bluegrass, particularly in the dormant season, limits the use of these community types for wildlife.

#### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

##### FORBS

##### DANDELION

(*Taraxacum officinale*) 15 6-37 100

##### WILD STRAWBERRY

(*Fragaria virginiana*) 3 0-21 61

##### CLOVER

(*Trifolium repens*) 15 0-52 74

##### YARROW

(*Achillea millefolium*) 7 0-15 96

##### GRACEFUL CINQUEFOIL

(*Potentilla gracilis*) 6 0-25 83

##### VEINY MEADOW RUE

(*Thalictrum venulosum*) 4 0-21 61

##### GRASSES

##### KENTUCKY BLUEGRASS

(*Poa pratensis*) 48 0-97 96

##### SLENDER WHEATGRASS

(*Agropyron trachycaulum*) 4 0-26 65

##### CREEPING RED FESCUE

(*Festuca rubra*) 3 0-26 30

##### TUFTED HAIRGRASS

(*Deschampsia cespitosa*) 1 0-4 22

#### ENVIRONMENTAL VARIABLES

##### MOISTURE REGIME:

MESIC

##### NUTRIENT REGIME:

MESOTROPHIC

##### ELEVATION:

1150-1600 M (1276 M)

##### SOIL DRAINAGE:

MODERATELY WELL

##### ASPECT:

VARIABLE

##### SLOPE:

0-35%

#### FORAGE PRODUCTION(KG/HA)

GRASS: 2206 (621-4319)

FORBS: 622 (153-2102)

SHRUBS: 150

TOTAL: 2837 (1014-4686)

#### SUGGESTED GRAZING CAPACITY

0.3 HA/AUM OR 0.7 ACRES/AUM

## UFC4. Kentucky bluegrass-Sedge/ Dandelion (*Poa pratensis*-*Carex* spp./ *Taraxacum officinale*)

n=25 This community type is similar to the Kentucky bluegrass/ clover-dandelion community type (UFC3), but it has not been grazed as heavily. There is still an abundance of native plant species such as veiny meadow rue, slender wheatgrass, tufted hairgrass and sedge, but there has been an increase in grazing resistant species, such as Kentucky bluegrass, dandelion and clover. If this community type is protected from grazing it will probably revert back to a tufted hairgrass-Kentucky bluegrass dominated type (UFC5) (Willoughby, 1992). Kentucky bluegrass, once established, appears to be a successful competitor.

These Kentucky bluegrass dominated community types are very productive, but they have lost two of the most advantageous species (tufted hairgrass and rough fescue). The forage quality of these native species is much better, particularly in the dormant season.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	2	0-6	48
PRICKLY ROSE ( <i>Rosa acicularis</i> )	2	0-19	16
<b>FORBS</b>			
DANDELION ( <i>Taraxacum officinale</i> )	24	1-53	100
YARROW ( <i>Achillea millefolium</i> )	9	1-25	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	6	0-14	84
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	11	0-41	80
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	14	0-40	88
<b>GRASSES</b>			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	36	0-85	96
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	5	0-29	64
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	6	0-21	72
SEDGE SPP. ( <i>Carex</i> spp.)	21	0-73	64
CREEPING RED FESCUE ( <i>Festuca rubra</i> )	1	0-5	12
ROUGH FESCUE ( <i>Festuca scabrella</i> )	4	0-12	60

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME

PERMESOTROPHIC

ELEVATION:

1150-1660 M (1447 M)

SOIL DRAINAGE:

MODERATELY WELL

### FORAGE PRODUCTION(KG/HA)

GRASS: 1915 (632-4304)

FORBS: 845 (264-3344)

SHRUBS: 102

TOTAL: 2767 (1491-4864)

<p>SUGGESTED GRAZING CAPACITY 0.3 HA/AUM OR 0.7 ACRES/AUM</p>
---

## UFC5. Tufted hairgrass-Kentucky bluegrass

(*Deschampsia cespitosa*-*Poa pratensis*)

n=4 This community type is similar to the other Kentucky bluegrass dominated community types, but grazing pressure has been lighter or it was heavy and then became more moderate because of reduced stocking rates or rotational grazing. Willoughby (1992), found that tufted hairgrass could compete with Kentucky bluegrass in the absence of grazing, but it appears that once Kentucky bluegrass is established it remains to form a stable community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	0-5	25
<b>FORBS</b>			
DANDELION ( <i>Taraxacum officinale</i> )	13	3-21	100
YARROW ( <i>Achillea millefolium</i> )	3	T-5	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	4	0-14	50
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	2	T-5	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	7	0-18	50
<b>GRASSES</b>			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	12	2-19	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	5	0-10	75
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	41	12-68	100
SEDGE SPP. ( <i>Carex spp.</i> )	9	0-17	100
ROUGH FESCUE ( <i>Festuca scabrella</i> )	2	0-5	75

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1300-1600 M (1400 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
NORTH

### FORAGE PRODUCTION(KG/HA)

GRASS: 3292  
FORBS: 1010  
TOTAL: 4302

<p>SUGGESTED GRAZING CAPACITY 0.2 HA/AUM OR 0.5 ACRES/AUM</p>
---

**UFC6. Sedge-Tufted hairgrass**  
(*Carex spp.-Deschampsia cespitosa*)

n=13 This community type was described at Harrison Flats in the Upper Clearwater River valley. It appears to represent a tufted hairgrass-sedge community that was heavily grazed in the past and now is rested and only lightly utilized. It appears that the heavy grazing pressure was not prolonged enough to allow Kentucky bluegrass invasion. It is also possible that Kentucky bluegrass is not predominant on this site because of lack of seed source in these isolated areas. It is likely, with continued protection from grazing, that this community type will succeed back to a modal tufted hairgrass-sedge dominated community type.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	2	0-6	69
<b>FORBS</b>			
DANDELION ( <i>Taraxacum officinale</i> )	7	0-22	69
YARROW ( <i>Achillea millefolium</i> )	11	0-41	85
STRAWBERRY ( <i>Fragaria virginiana</i> )	5	0-10	54
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	10	0-38	62
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	5	0-26	46
<b>GRASSES</b>			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	1	0-5	23
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	2	0-13	31
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	20	0-46	92
SEDGE ( <i>Carex spp.</i> )	63	0-93	100
ROUGH FESCUE ( <i>Festuca scabrella</i> )	7	0-19	62
BALTIC RUSH ( <i>Juncus balticus</i> )	15	0-58	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1529-1829 M (1781 M)  
SOIL DRAINAGE:  
IMPERFECTLY  
ASPECT:  
VARIABLE  
SLOPE:  
0%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1020(684-1510)  
FORBS: 409 (150-891)  
SHRUBS: 322  
TOTAL: 1468 (932-1962)

**SUGGESTED GRAZING CAPACITY**  
0.6 HA/AUM OR 1.4 ACRES/AUM

## UFC7. Creeping red fescue-Clover

(*Festuca rubra*-*Trifolium spp.*)

n=23 This community type was described at lower elevations, adjacent to farms and ranches in the Upper Foothills subregion. This community represents native communities that have been disturbed and planted to creeping red fescue. These include pipelines, roadsides and cultivated pastures. Lane et al. (2000), felt this community type developed when a site which was seeded to creeping red fescue-timothy-clover and received low levels of use. Creeping red fescue spreads throughout the site by creeping rhizomes and chokes out the timothy by forming a dense matt of litter. This community type is normally considered to be in good or excellent condition.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

FORBS			
DAÑDELION ( <i>Taraxacum officinale</i> )	7	0-21	88
YARROW ( <i>Achillea millefolium</i> )	2	0-7	1
STRAWBERRY ( <i>Fragaria virginiana</i> )	2	0-12	63
CLOVER SPP. ( <i>Trifolium spp</i> )	22	0-49	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	1	0-7	29
GRASSES			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	18	0-58	83
CREEPING RED FESCUE ( <i>Festuca rubra</i> )	37	6-87	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	1	0-4	25
SEDGE SPP. ( <i>Carex spp.</i> )	2	0-11	58
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	2	0-17	29
TIMOTHY ( <i>Phleum pratense</i> )	5	0-31	71

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME  
MESIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1212-1880 M (1464 M)  
SOIL DRAINAGE:  
WELL  
ASPECT:  
VARIABLE  
SLOPE:  
0-40%

### FORAGE PRODUCTION(KG/HA)

GRASS: 1789 (332-4894)  
FORBS: 302 (20-1114)  
SHRUBS: 384  
TOTAL: 2009 (404-5054)

SUGGESTED GRAZING CAPACITY 0.5 HA/AUM OR 1.0 ACRES/AUM
---

**UFC8. Kentucky bluegrass-Timothy/ Meadow rue**  
*(Poa pratensis-Phleum pratense/ Thalictrum venulosum)*

n=4 This community type represents the grazed and disturbed community of the cow parsnip-meadow rue/fringed brome community (UFA14). The high productivity and open nature of this community make it extremely attractive to domestic livestock. Heavy to moderate grazing pressure causes cow parsnip, veiny meadow rue and fringed brome to decrease and allows Kentucky bluegrass, timothy and dandelion to increase.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp.</i> )	2	0-5	75
<b>FORBS</b>			
DANDELION			
( <i>Taraxacum officinale</i> )	13	2-35	100
COW PARSNIP			
( <i>Heracleum lanatum</i> )	7	0-23	50
STRAWBERRY			
( <i>Fragaria virginiana</i> )	1	0-1	50
CLOVER			
( <i>Trifolium repens,</i> <i>T. hybridum</i> )	5	0-20	25
VEINY MEADOW RUE			
( <i>Thalictrum venulosum</i> )	7	0-24	50
<b>GRASSES</b>			
KENTUCKY BLUEGRASS			
( <i>Poa pratensis</i> )	22	0-33	75
SLENDER WHEATGRASS			
( <i>Agropyron trachycaulum</i> )	7	0-16	75
SMOOTH BROME			
( <i>Bromus inermis</i> )	2	0-7	25
TIMOTHY			
( <i>Phleum pratense</i> )	17	11-25	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1060-15201M (1330 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-6%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 2834 (308-6322)  
 FORBS: 1469 (210-2830)  
 SHRUBS: 42  
 TOTAL: 4313 (560-6942)

**SUGGESTED GRAZING CAPACITY**  
 0.3 HA/AUM OR 0.7 ACRES/AUM

**UFC9. Purple oatgrass-Rough fescue**  
*(Schizachne purpurascens -Festuca scabrella)*

n=1 This community type was described in the Ghost area west of Calgary. It appears to represent a rough fescue, hairy wildrye dominated grassland that has undergone heavy grazing pressure. Willoughby (1995) has found that purple oatgrass will increase with increased grazing pressure on nutrient poor soils in the Lower Foothills subregion. Willoughby (2000) also described a purple oatgrass-california oatgrass dominated community type on saline soils in the Dry Mixedwood subregion. It is possible that this community type maybe associated with a saline seepage area which favours the growth of purple oatgrass.

This community type is fairly productive, but the majority of the production is coming from purple oatgrass which is only moderately palatable to livestock. This community type should probably be rated as secondary range.

**PLANT COMPOSITION**

**CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL			
<i>(Potentilla fruticosa)</i>	6	-	100
<b>FORBS</b>			
THREE FLOWERED AVENS			
<i>(Geum triflorum)</i>	4	-	100
GRACEFUL CINQUEFOIL			
<i>(Potentilla gracilis)</i>	3	-	100
YARROW			
<i>(Achillia millifolium)</i>	3	-	100
SMOOTH ASTER			
<i>(Aster laevis)</i>	2	-	100
<b>GRASSES</b>			
PURPLE OAT GRASS			
<i>(Schizachne purpurascens)</i>	15	-	100
SLENDER WHEATGRASS			
<i>(Agropyron trachycaulum)</i>	10	-	100
ROUGH FESCUE			
<i>(Festuca scabrella)</i>	7	-	100
PRESL SEDGE			
<i>(Carex preslii)</i>	5	-	100
JUNEGRASS			
<i>(Koeleria macrantha)</i>	1	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBMESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1460  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTHERLY  
 SLOPE:  
 25%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 2052  
 FORBS: 362  
 SHRUBS: 384  
 TOTAL: 2798

**SUGGESTED GRAZING CAPACITY**  
 0.3 Ha/AUM or 0.7 Ac/AUM

**UFC10. Willow/ Kentucky bluegrass**  
*(Salix spp./Poa pratensis)*

n=2 This community type represents the grazed and disturbed community of the willow/ tufted hairgrass-sedge community type (UFB3). The high productivity and open nature of this community make it extremely attractive to domestic livestock. Heavy to moderate grazing pressure causes the native plant species to decrease and allows Kentucky bluegrass and dandelion to increase.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW			
<i>(Salix spp.)</i>	22	19-25	100
<b>FORBS</b>			
DANDELION			
<i>(Taraxacum officinale)</i>	11	1-21	100
YARROW			
<i>(Achillea millefolium)</i>	7	5-7	100
STRAWBERRY			
<i>(Fragaria virginiana)</i>	2	T-4	100
WHITE CLOVER			
<i>(Trifolium repens)</i>	5	0-10	50
TALL LUNGWORD			
<i>(Mertensia paniculata)</i>	5	0-10	50
<b>GRASSES</b>			
KENTUCKY BLUEGRASS			
<i>(Poa pratensis)</i>	12	10-13	100
SLENDER WHEATGRASS			
<i>(Agropyron trachycaulum)</i>	9	5-12	100
TUFTED HAIRGRASS			
<i>(Deschampsia cespitosa)</i>	8	0-15	50
SHEEP FESCUE			
<i>(Festuca saximontana)</i>	5	0-10	50

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1370-1667 M (1499 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-25%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 1466 (880-2052)  
 FORBS: 339 (316-332)  
 SHRUBS: 384  
 TOTAL: 1997 (1196-2798)

<p><b>SUGGESTED GRAZING CAPACITY</b>                  0.5 HA/AUM OR 1.1 ACRES/AUM</p>
---

**UFC11. Sedge-Slender wheatgrass-Rough fescue**  
*(Carex spp.-Agropyron trachycaulum-Festuca scabrella)*

n=29 This community type represents the grazed transects at the McCue Creek, Yara Creek and Upper James River rangeland reference areas over 30 years from the 1960's to the early 1980's. The continued grazing pressure since the 1980's has allowed Kentucky bluegrass to invade onto these sites to form a Kentucky bluegrass-Sedge dominated community type (Willoughby 2000). In the 1960's when these sites were protected from grazing the plant community succeeded back to a rough fescue-hairy wildrye dominated community.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa.)</i>	2	0-10	62
<b>FORBS</b>			
DANDELION <i>(Taraxacum officinale)</i>	3	0-13	90
YARROW <i>(Achillea millefolium)</i>	3	0-7	97
STRAWBERRY <i>(Fragaria virginiana)</i>	2	0-18	45
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	3	0-8	79
AMERICAN VETCH <i>(Vicia americana)</i>	3	0-16	93
<b>GRASSES</b>			
KENTUCKY BLUEGRASS <i>(Poa pratensis)</i>	3	0-16	52
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	9	1-19	100
ROUGH FESCUE <i>(Festuca scabrella)</i>	8	1-32	100
SEDGE SPP. <i>(Carex spp.)</i>	10	0-31	69
HAIRY WILDRYE <i>(Elymus innovatus)</i>	3	0-20	41

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
 NUTRIENT REGIME:  
MESOTROPHIC  
 ELEVATION:  
1470-1660(1543M)  
 SOIL DRAINAGE:  
WELL  
 ASPECT:  
SOUTHERLY  
 SLOPE:  
2-22%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 802  
 FORBS: 322  
 SHRUBS: 18  
 TOTAL: 1207

**SUGGESTED GRAZING CAPACITY**  
 1.1 HA/AUM OR 2.5 ACRES/AUM



**UPPER FOOTHILLS SUBREGION**  
**DECIDUOUS COMMUNITY TYPES**



**Figure 5. Aspen/Marsh reedgrass community type on south facing slopes in the Upper Foothills subregion**



### Deciduous community types

The six deciduous community types described in the Upper Foothills subregion are outlined in Table 4. Deciduous types are rare in this subregion. The cool climate severely restricts the growth of deciduous tree species (Strong and Leggat 1992). As a result, aspen and balsam poplar are generally found on south facing slopes where the increased insolation permits colonization.

The Pb-Sw/ Willow/ Yellow Mtn. avens community type (UFD2) is representative of the gravelly floodplains adjacent to rivers and streams. The Aw/ Rose/ Bearberry, Aw/ Rose/ Hairy wildrye and Aw/ Buffaloberry/ Hairy wildrye community types are found on dry south facing slopes throughout the region. The Aw/ Buffaloberry/ Hairy wildrye community type appears to be successionaly more advanced, with slightly acidic soils, than the Aw/ Rose/ Hairy wildrye community type. The Aw/ Marsh reedgrass type is slightly moister than the other aspen community types found on south facing slopes in the Upper Foothills subregion and the Pb/ Willow/ Horsetail was described on the river floodplain adjacent to the Wildhay river.

**Key to Deciduous Community Types**

- 1. balsam poplar dominates the overstory. .... 2  
Drier sites with Aspen dominating the overstory ..... 3
- 2. Low nutrient, dry gravelly river floodplains .. UFD2 Balsam poplar-White spruce/ Willow/ Yellow mtn. Avens  
Moist, moderately well drained soils ..... UFD6 Balsam poplar/ Willow/ horsetail
- 3. Dry , south and west facing slopes ..... 4  
Moister richer sites with Marsh reedgrass dominated understory ..... UFD5 Aspen/ Marsh reedgrass
- 4. Bearberry or Buffaloberry dominate understory. .... 5  
Grass and forbs dominate the understory of this dry, sunny site ..... UFD3 Aspen/ Rose/ Hairy wildrye
- 5. Dry site conditions on steep south facing slope ..... UFD1 Aspen/ Rose/ Bearberry  
Lower, south facing slopes and lower nutrient soils (high pH) ..... UFD4 Aspen/ Buffaloberry/ Hairy wildrye

Table 4. Forage production (kg/ha) for the deciduous community types and ecosite phases within the Upper Foothills subregion

Community number	Community type	Productivity(kg/ha)			Moisture	Drainage	Carrying capacity(Ha/AUM)	
		Grass	Forb	Shrub				Total
<b>c2</b>	<b>Ecosite phase hairy wildrye Aw</b>	350	372	149	871	Submesic	Well	1.4
UFD1	Aw/Rose/Bearberry	450	300	114	864	Submesic	Well	2.1
UFD3	Aw/Rose/Hairy wildrye	200	467	134	800	Mesic	Mod. Well	1.1
UFD4	Aw/Bufaloberry/Hairy wildrye	400	350	200	950	Submesic	Well	1.0
<b>c5</b>	<b>Ecosite phase yellow mtn avens</b>	62	316	230	608	Mesic	Imperfectly	3.0
UFD2	Pb-Sw/Willow/Yellow Mtn. Avens	62	316	230	608	Mesic	Imperfectly	3.0
<b>f2</b>	<b>Ecosite phase bracted honeysuckle Aw</b>	206	776	110	1092	Subhygric	Mod. Well	1.6
UFD5	Aw/Marsh reedgrass	206	776	110	1092	Subhygric	Mod. Well	1.6
<b>j2</b>	<b>Ecosite phase horsetail Pb</b>	50	550	150	750	Subhygric	Mod. Well	1.2
UFD6	Pb/Willow/Horsetail	50	550	150	750	Subhygric	Mod. Well	1.2

**UFD1. Aspen/ Rose/ Bearberry**  
*(Populus tremuloides/ Rosa acicularis/ Arctostaphylos uva-ursi)*

n=1 This community type was described on the steep south facing slope above Rough Creek, west of Rocky Mountain House. The drier site conditions favour the growth of bearberry. This community type probably represents an earlier successional phase of the PI/ bearberry community type described by Beckingham et al (1996). The forage productivity of this community type is only moderate, but the openness of the stand makes it accessible for livestock. This community type would be rated as secondary range.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>TREES</b>			
ASPEN			
<i>(Populus tremuloides)</i>	47	-	100
<b>SHRUBS</b>			
PRICKLY ROSE			
<i>(Rosa acicularis)</i>	22	-	100
BEARBERRY			
<i>(Arctostaphylos uva-ursi)</i>	16	-	100
<b>FORBS</b>			
STRAWBERRY			
<i>(Fragaria virginiana)</i>	7	-	100
LINDLEY'S ASTER			
<i>(Aster ciliolatus)</i>	1	-	100
FIREWEED			
<i>(Epilobium angustifolium)</i>	11	-	100
YARROW			
<i>(Achillea millefolium)</i>	3	-	100
<b>GRASSES</b>			
SLENDER WHEATGRASS			
<i>(Agropyron trachycaulum)</i>	5	-	100
FRINGED BROME			
<i>(Bromus ciliatus)</i>	3	-	100
ROUGH FESCUE			
<i>(Festuca scabrella)</i>	3	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1215 M  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTH-EAST  
 SLOPE:  
 30%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 450  
 FORBS: 300  
 SHRUBS: 114  
 TOTAL: 864

SUGGESTED GRAZING CAPACITY  
 2.1 HA/AUM OR 5.1 ACRES/AUM

**UFD2. Balsam poplar/ Willow/ Yellow Mountain Avens**  
*(Populus balsamifera/ Salix spp./ Dryas drummondii)*

n=1 This community type is common throughout the Upper Foothills subregion on gravelly floodplains along rivers and streams. It is similar to the bearberry/ slender wheatgrass community (UFA10), but it is successional more advanced. This type is dominated by balsam poplar with an understory of spruce in the later successional stages. This particular stand was fairly young with the tree canopy being less than 5 m tall. Yellow mountain avens is a common pioneer species on gravelly river bars and rocky slopes up into the alpine tundra (MacKinnon et al., 1992). As this community succeeds towards a mature forest, yellow mountain avens will undoubtedly decline in cover.

The forage production on this community type is very low. The poor nutrient status of the soil limits the growth of grasses, forbs and shrubs. As a result, this community type would be rated as non-use range.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
BALSAM POPLAR <i>(Populus balsamifera)</i>	7	-	100
WHITE SPRUCE <i>(Picea glauca)</i>	5	-	100
<b>SHRUBS</b>			
WILLOW <i>(Salix spp.)</i>	13	-	100
BUFFALOBERRY <i>(Shepherdia canadensis)</i>	9	-	100
YELLOW MOUNTAIN AVENS <i>(Dryas drummondii)</i>	16	-	100
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	3	-	100
<b>FORBS</b>			
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	1	-	100
ALPINE HEDYSARUM <i>(Hedysarum alpinum)</i>	11	-	100
SCOURING RUSH <i>(Equisetum scirpoides)</i>	11	-	100
ALPINE MILKVETCH <i>(Astragalus alpinus)</i>	4	-	100
<b>GRASSES</b>			
BLUNT SEDGE <i>(Carex obtusata)</i>	2	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1524 M  
SOIL DRAINAGE:  
IMPERFECTLY

**FORAGE PRODUCTION(KG/HA)**

GRASS: 62  
FORBS: 316  
SHRUBS: 230  
TOTAL: 608

SUGGESTED GRAZING CAPACITY  
3 HA/AUM OR 7 ACRES/AUM

**UFD3. Aspen/ Rose/ Hairy wildrye**  
*(Populus tremuloides/ Rosa acicularis/ Elymus innovatus)*

n=10 This community type is typical of aspen forest types found throughout the Upper Foothills subregion on south facing slopes. The dry site conditions and high solar insolation favours the growth of grasses and forbs rather than shrubs. The canopy cover of aspen is also noticeably lower on this community type. This community type is similar to the Aw/ buffaloberry/ hairy wildrye community (UFD4) described in Willmore Wilderness Park, but the absence of buffaloberry distinguishes this type from the Willmore type. This community is moderately productive for domestic livestock. This community would be rated as secondary range.

**PLANT COMPOSITION** **CANOPY COVER(%)**

MEAN RANGE CONST.

<b>TREES</b>			
<b>ASPEN</b>			
<i>(Populus tremuloides)</i>	41	10-72	100
<b>SHRUBS</b>			
<b>PRICKLY ROSE</b>			
<i>(Rosa acicularis)</i>	1	0-7	60
<b>WILLOW</b>			
<i>(Salix spp.)</i>	1	0-10	40
<b>FORBS</b>			
<b>LINDLEY'S ASTER</b>			
<i>(Aster ciliolatus)</i>	4	0-18	40
<b>WILD STRAWBERRY</b>			
<i>(Fragaria virginiana)</i>	13	1-33	100
<b>TALL LUNGWORT</b>			
<i>(Mertensia paniculata)</i>	4	0-12	100
<b>CREAM COLORED VETCHLING</b>			
<i>(Lathyrus ochroleucus)</i>	3	1-7	100
<b>VEINY MEADOW RUE</b>			
<i>(Thalictrum venulosum)</i>	3	0-9	80
<b>GRASSES</b>			
<b>HAIRY WILD RYE</b>			
<i>(Elymus innovatus)</i>	20	0-62	90
<b>PURPLE OATGRASS</b>			
<i>(Schizachne purpurascens)</i> 4		0-20	40
<b>SLENDER WHEATGRASS</b>			
<i>(Agropyron trachycaulum)</i> 4		0-26	20

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME  
 MESIC  
 NUTRIENT REGIME  
 MESOTROPHIC  
 ELEVATION:  
 1345-1515 M (1457 M)  
 SOIL DRAINAGE (MEAN):  
 MODERATELY WELL  
 ASPECT:  
 SOUTH TO SOUTHWEST  
 SLOPE:  
 0-35%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 200  
 FORBS: 467 (150-1000)  
 SHRUBS: 134 (50-300)  
 TOTAL: 800 (401-1500)

**SUGGESTED GRAZING CAPACITY**  
 1.1 HA/AUM OR 2.5 AC/AUM

**UFD4. Aspen/ Buffaloberry/ Hairy wildrye**  
*(Populus tremuloides/ Shepherdia canadensis/ Elymus innovatus)*

n=3 This community type was described along lower, south facing slopes and river terraces throughout Willmore Wilderness Park and areas west of Hinton. Bork (1994), found this community type. to be uncommon throughout Willmore, but pockets of this type were found along the Smoky, Sulphur and Sheep rivers on the North side of the Park. Bork felt that frequent disturbance and/ or arid conditions resulted in the aspen dominated overstory. He felt if left undisturbed, the community type would eventually succeed to a coniferous forest. This community type is very similar to the Aw/ buffaloberry type described by Youngblood (1993) in Alaska and the Aw/ rose/ hairy wildrye community type (UFD3) previously described near Rocky Mtn. House. The presence of buffaloberry distinguishes this northern type from the more southern rose type. The presence of buffaloberry may indicate a higher pH and lower nutrient status. Beckingham (1994), described Aw/ buffaloberry stands on lower pH sites.

This community type provides a good forage base for domestic livestock. In the Upper foothills, this community type is often located in close proximity to the trails and camps used by outfitters and recreationalists.

**PLANT COMPOSITION** **CANOPY COVER(%)**

MEAN RANGE CONST.

**TREES**

**ASPEN**

*(Populus tremuloides)* 34 24-52 100

**WHITE SPRUCE**

*(Picea glauca)* 4 0-11 33

**SHRUBS**

**BUFFALOBERRY**

*(Shepherdia canadensis)* 14 10-18 100

**PRICKLY ROSE**

*(Rosa acicularis)* 7 1-7 100

**WILLOW**

*(Salix spp.)* 17 5-36 100

**TWIN-FLOWER**

*(Linnaea borealis)* 1 0-4 33

**BUNCHBERRY**

*(Cornus canadensis)* 2 0-7 33

**FORBS**

**FIREWEED**

*(Epilobium angustifolium)* 3 1-5 100

**SHOWY ASTER**

*(Aster conspicuus)* 1 0-4 33

**STRAWBERRY**

*(Fragaria virginiana)* 9 3-19 100

**GRASSES**

**HAIRY WILD RYE**

*(Elymus innovatus)* 24 14-34 100

**MARSH REEDGRASS**

*(C.alamagrostis canadensis)* 3 0-5 50

**ENVIRONMENTAL VARIABLES**

**MOISTURE REGIME:**

MESIC

**NUTRIENT REGIME:**

MESOTROPHIC

**ELEVATION:**

914-1500 M (957 M)

**SOIL DRAINAGE:**

WELL

**ASPECT:**

SOUTH

**SLOPE:**

0-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 400

FORBS: 350

SHRUBS: 200

TOTAL: 950

**SUGGESTED GRAZING CAPACITY**

1 HA/AUM OR 2.1 ACRES/AUM

**UFD5. Aspen/ Marsh reedgrass**  
(*Populus tremuloides/ Calamagrostis canadensis*)

n=4 This community type was described on a south facing slope in the Solomon valley west of Hinton and observed near Fall Creek and Upper James west of Rocky Mountain House. This community type is scattered throughout the valleys in small isolated areas. It appears to have a slightly higher moisture regime than the bearberry, hairy wildrye and buffalobery dominated community types previously described. The dominance of marsh reedgrass indicates that some nutrient rich seepage occurs at some point in the growing season. This community type was located adjacent to Kentucky bluegrass-timothy dominated meadows (UFC8). As a result, this aspen dominated community type was extensively utilized by livestock.

**PLANT COMPOSITION**

CANOPY COVER(%)  
MEAN RANGE CONST.

<b>TREES</b>			
ASPEN			
( <i>Populus tremuloides</i> )	29	14-45	100
BALSAM POPLAR			
( <i>Populus balsamifera</i> )	3	0-8	75
WHITE SPRUCE			
( <i>Picea glauca</i> )	7	0-13	75
<b>SHRUBS</b>			
PRICKLY ROSE			
( <i>Rosa acicularis</i> )	1	0-3	75
WILLOW			
( <i>Salix spp.</i> )	3	0-8	50
<b>FORBS</b>			
CANADA VIOLET			
( <i>Viola canadensis</i> )	4	0-17	50
PEAVINE			
( <i>Lathyrus ochroleucus</i> )	3	0-5	75
TALL LUNGWORT			
( <i>Mertensia paniculata</i> )	3	T-6	100
LINDLEY'S ASTER			
( <i>Aster ciliolatus</i> )	5	0-13	50
STRAWBERRY			
( <i>Fragaria virginiana</i> )	4	1-9	100
COW PARNISP			
( <i>Heracleum lanatum</i> )	3	0-10	50
<b>GRASSES</b>			
HAIRY WILD RYE			
( <i>Elymus innovatus</i> )	6	3-10	100
MARSH REEDGRASS			
( <i>Calamagrostis canadensis</i> )	14	4-20	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1450-1500 M (1477 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
WEST  
SLOPE:  
3-18%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 206 (110-301)  
FORBS: 776 (350-1202)  
SHRUBS: 110 (100-120)  
TOTAL: 1092 (751-1432)

**SUGGESTED GRAZING CAPACITY**  
1.6 HA/AUM OR 3.8 AC/AUM

**UFD6. Balsam poplar/ Willow/ Horsetail**  
*(Populus balsamifera/ Salix spp/ Equisetum arvense)*

n=1 This community type was described on the flood plain of the Wildhay River northwest of Hinton. This community is not common in the Upper Foothills subregion and likely represents the continued succession of a willow/ horsetail dominated community type (UFB12). Continued succession in the absence of disturbance will likely lead to the development of a Sw/ horsetail dominated community type (UFE6).

This community type is being used by livestock because of its close proximity to a right of way that had been seeded to Creeping red fescue and clover. When in close proximity to primary range areas this community type should be rated as secondary range.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

<b>TREES</b>			
ASPEN			
<i>(Populus tremuloides)</i>	5	-	100
<b>BALSAM POPLAR</b>			
<i>(Populus balsamifera)</i>	35	-	100
<b>WHITE SPRUCE</b>			
<i>(Picea glauca)</i>	3	-	100
<b>SHRUBS</b>			
<b>PRICKLY ROSE</b>			
<i>(Rosa acicularis)</i>	3	-	100
<b>BEAKED WILLOW</b>			
<i>(Salix bebbiana)</i>	50	-	100
<b>FORBS</b>			
<b>HORSETAIL</b>			
<i>(Equisetum arvense)</i>	12	-	100
<b>SCOURING RUSH</b>			
<i>(Equisetum scirpoides)</i>	9	-	100
<b>TALL LUNGWORT</b>			
<i>(Mertensia paniculata)</i>	3	-	100
<b>LINDLEY'S ASTER</b>			
<i>(Aster ciliolatus)</i>	4	-	100
<b>STRAWBERRY</b>			
<i>(Fragaria virginiana)</i>	7	-	100
<b>RED CLOVER</b>			
<i>(Trifolium pratense)</i>	4	-	100
<b>GRASSES</b>			
<b>HAIRY WILD RYE</b>			
<i>(Elymus innovatus)</i>	1	-	100
<b>MARSH REEDGRASS</b>			
<i>(Calamagrostis canadensis)</i>	1	-	100
<b>KENTUCKY BLUEGRASS</b>			
<i>(Poa pratensis)</i>	1	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1500 M  
 SOIL DRAINAGE:  
 MODERATELY WELL

**FORAGE PRODUCTION(KG/HA)**

GRASS: 50  
 FORBS: 550  
 SHRUBS: 150  
 TOTAL: 750

<p><b>SUGGESTED GRAZING CAPACITY</b>          1.2 HA/AUM OR 2.7 AC/AUM</p>
--



## Conifer community types

Lodgepole pine dominates the overstory vegetation of the mesic reference sites in the Upper Foothills subregion. Typical forests are represented by the Pl/ bog cranberry (UFE1) and the Pl/ marsh reedgrass c.t. (UFE4) community types. Secondary succession is by white spruce and leads to the formation of the Pl-Sw/ bunchberry c.t. (UFE2). Continued succession on wetter sites, in the absence of disturbance, leads to a Sw/ horsetail/ moss dominated c.t. (UFE6) and to the Sw/ moss (UFE10) dominated community on more mesic sites.

Wetter, subhygric sites can be dominated by lodgepole pine, black spruce or white spruce. Many of these sites have a predominant willow understory (Pl/ willow/ moss (UFE3) or Sw/ willow(UFE7)). These types appear to represent continued succession from the native shrub and grassland community types. Succession in the absence of disturbance on these sites will be to white spruce. The Sw/ willow c.t. (UFE7) appears to be typical of a climax forest on these subhygric sites.

Black spruce dominates poorly drained depressional areas (Sb/ willow (UFE5)). These sites have a high water table throughout most of the year. Organic accumulations are a common result of the poor drainage conditions and low oxygen availability (Strong and Leggat, 1992).

Dry, south facing slopes are typically dominated by deciduous aspen forests with succession to a Sw/ bearberry (UFE8) and Pl/ bearberry/ hairy wildrye dominated community types (Beckingham et al., 1996). A Sw/ juniper (UFE9) c.t. was described on fine-textured, calcareous loess deposits, with high pH's near Brule lake. These deposits blow out of the Athabasca river valley from Jasper National Park.

The conifer forest types are generally unsuitable for livestock grazing and are typically rated as non-use. The ten coniferous community types described in the Upper Foothills subregion are outlined in Table 5. A more complete description of coniferous c.t. can be found in Beckingham et al. (1996).

**Key to Coniferous Community Types**

1. Lodgepole pine dominated community ..... 2  
 White or black spruce dominated community ..... 5
2. The community is supported by well drained, coarse soil. It is often located on south facing slopes and exhibits a structured understory of grass, forb and shrub species ..... 3  
 The community is located on less developed, wet soils which support a willow understory ..... UFE3 Lodgepole pine/ Willow/ Moss
3. The community is succeeding to white spruce in undisturbed areas. Moss cover increases with canopy closure and grass, forb, and shrub species decline ..... UFE2 Lodgepole pine-White spruce/ Bunchberry  
 The understory is dominated by shrubs, forbs or grasses ..... 4
4. Shrubs dominate the understory on these well drained, south slopes ..... UFE1 Lodgepole pine/ Bog cranberry  
 Forbs and grasses dominate the understory ..... UFE4 Lodgepole pine/ Marsh reedgrass
5. Dry site conditions ..... 6  
 Wet site conditions ..... 7
6. Poor nutrient regime, bearberry dominates the understory, wind ..... UFE8 White spruce/ Bearberry  
 Fine textured calcareous loess with a high pH ..... UFE9 White spruce/ Juniper-Buffaloberry
7. Drainage is poor, willow dominates the understory ..... 8  
 Mesic, moss dominates the understory ..... 9
8. White spruce dominated overstory, wetter, low light ..... UFE7 White spruce/ Willow  
 Black spruce dominated overstory, wet, often saturated soils ..... UFE5 Black spruce/ Willow
9. Wet soils, canopy is closing, promoting moss and loss of forb etc. .... UFE6 White spruce/ Horsetail/ Moss  
 Closed canopy, successional mature ..... UFE10 White spruce/ Moss

Table 5. Conifer community types of the Upper Foothills subregion

Community name	Community type	Productivity(kg/Ha)		Moisture	Drainage	Carrying capacity (Ha/AUM)		
		Grass	Forb					
<b>c4 Ecosite phase hairy wildrye Sw</b>		297	176	181	517	Mesic	Mod. Well	Non-use
UFE8.	White spruce/Bearberry				400	Mesic	Mod. Well	Non-Use
UFE9.	White spruce/Juniper-Buffaloberry	297	176	181	634	Mesic	Well	Non-Use
<b>e1 Ecosite phase t. bilberry/arnica P1</b>		62	316	92	478	Mesic	Well	Non-use
UFE1.	Lodgepole pine/Bog cranberry	62	316	92	271	Mesic	Well	Non-use
UFE2.	Lodgepole pine-White spruce/ Bunchberry				361	Mesic	Well	Non-Use
UFE4.	Lodgepole pine/Marsh reedgrass				801	Mesic	Well	Non-Use
<b>e3 Ecosite phase t. bilberry/arnica Sw</b>		78	96	160	332	Mesic	Well	Non-use
UFE10	White spruce/Moss	78	96	160	332	Mesic	Well	Non-Use
<b>f1 Ecosite phase bracted honeysuckle P1</b>		478	192	252	845	Subhygric	Mod. Well	Non-use
UFE3.	Lodgepole pine/Willow/Moss	478	192	252	845	Subhygric	Mod. Well	Non-Use
<b>j1 Ecosite phase horsetail Sw</b>		83	223	98	359	Subhygric	Mod. Well	Non-use
UFE6.	White spruce/Horsetail/Moss	83	223	98	418	Subhygric	Mod. Well	Non-Use
UFE7.	White spruce/Willow				300*	Subhygric	Mod. Well	Non-Use
<b>k1 Ecosite phase treed bog</b>		89	166	130	385	Hygric	Imperfectly	Non-use
UFE5.	Black spruce/Willow	89	166	130	385	Hygric	Imperfectly	Non-Use

\* Estimate

## UFE1. Lodgepole pine/ Bog cranberry

(*Pinus contorta*/ *Vaccinium vitis-idaea*)

n=8 This community type is common on dry, coarse textured, well drained sites throughout the Upper Foothills subregion and is part of the suberic/ poor ecosite described by Beckingham et al. (1996). These sites are generally located on slopes with southerly aspects. This community type is very similar to the PI/ hairy wildrye/ bunchberry community type described by Lane et al. (2000) in the Lower Foothills subregion, and the PI-Sw/ low bush cranberry/ twinflower type described by Beckingham (1994) in the Upper Foothills subregion, but this community type appears to be drier with a poorer nutrient regime. Beckingham (1994), felt that white spruce would eventually dominate the canopy of this community type.

Generally, this community type is not useful for domestic livestock grazing because it does not produce good quality forage.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### TREES

LOGEPOLE PINE  
(*Pinus contorta*) 35 20-50 100

WHITE SPRUCE  
(*Picea glauca*) 7 0-20 50

#### UNDERSTORY TREES

WHITE SPRUCE  
(*Picea glauca*) 2 0-15 25

#### SHRUBS

BOG CRANBERRY  
(*Vaccinium vitis-idaea*) 22 9-57 100

LABRADOR TEA  
(*Ledum groenlandicum*) 4 0-18 63

DEWBERRY  
(*Rubus Pubescens*) 2 0-14 13

BEARBERRY  
(*Arctostaphylos uva-ursi*) 1 0-9 13

BUNCHBERRY  
(*Cornus canadensis*) 5 0-14 88

TWINFLOWER  
(*Linnaea borealis*) 6 0-21 88

#### GRASSES

HAIRY WILDRYE  
(*Elymus innovatus*) 6 0-18 88

MOSS 63 27-86 100

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

MESIC

#### NUTRIENT REGIME:

SUBMESOTROPHIC

#### ELEVATION:

1091-1475 M (1354 M)

#### SOIL DRAINAGE:

WELL

#### ASPECT:

VARIABLE

#### SLOPE:

0-15%

### FORAGE PRODUCTION (KG/HA)

GRASS: 62

FORBS: 316

SHRUBS: 92

TOTAL 271 (89-470)

SUGGESTED GRAZING CAPACITY  
NON-USE

**UFE2. Lodgepole pine-White spruce/ Bunchberry**  
*(Pinus contorta-Picea glauca/ Cornus canadensis)*

n=5 This community type represents the modal type on mesic/ mesotrophic sites throughout the Upper Foothills subregion and may be transitional to the Lower Foothills subregion if aspen occurs in the stand. Strong (1992), found that lodgepole pine dominated the reference sites in this subregion with white spruce succession occurring on undisturbed areas. Beckingham (1994), described a similar community type (P1-Sw/ low bush cranberry/ twinflower) and felt that white spruce and balsam fir will eventually dominate the canopy. The change in canopy dominance will lead to a decline in understory cover of shrubs and forbs. As succession occurs, moss cover will increase.

This community type would be rated as non-use range for domestic livestock. There is little forage that

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

	MEAN	RANGE	CONST.
<b>TREES</b>			
LOGEPOLE PINE <i>(Pinus contorta)</i>	37	30-45	100
WHITE SPRUCE <i>(Picea glauca)</i>	21	0-35	80
<b>SHRUBS</b>			
DWARF BILBERRY <i>(Vaccinium caespitosum)</i>	3	0-7	80
BOG CRANBERRY <i>(Vaccinium vitis-idaea)</i>	3	0-5	80
WILLOW <i>(Salix spp.)</i>	2	0-5	60
BUNCHBERRY <i>(Cornus canadensis)</i>	21	2-39	100
TWIN-FLOWER <i>(Linnaea borealis)</i>	2	0-5	100
<b>FORBS</b>			
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	1	0-3	60
<b>GRASSES</b>			
HAIRY WILD RYE <i>(Elymus innovatus)</i>	5	0-12	100
MARSH REEDGRASS <i>(Calamagrostis canadensis)</i>	1	0-1	80
MOSS	59	36-76	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1091-1500 M (1368 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTH  
 SLOPE:  
 0-15%

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 361 (288-496)

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFE3. Lodgepole pine/ Willow/ Moss**  
(*Pinus contorta/ Salix spp./ Moss spp.*)

n=3 This community type is very similar to the other lodgepole pine dominated community types, but it is found on wetter soils that lack development. This community type is slightly drier than the Pl-Sb/ labrador tea-whortleberry/ bunchberry/ feather moss type described by Beckingham (1994) and the Sb/ willow dominated community type (UFE5) described in this guide. Herbaceous plants are scarce in the understory of this community type. As a result, there is little forage for domestic livestock and this community would be rated non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

	MEAN	RANGE	CONST.
<b>TREES</b>			
LODGEPOLE PINE ( <i>Pinus contorta</i> )	32	25-40	100
WHITE SPRUCE ( <i>Picea glauca</i> )	13	5-30	100
<b>SHRUBS</b>			
WILLOW ( <i>Salix spp.</i> )	23	13-34	100
BUNCHBERRY ( <i>Cornus canadensis</i> )	4	1-6	100
TWIN-FLOWER ( <i>Linnaea borealis</i> )	1	0-3	33
<b>FORBS</b>			
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	3	1-4	100
PALMATE LEAVED COLTSFOOT ( <i>Petasites palmatus</i> )	1	T	100
<b>GRASSES</b>			
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	4	T-7	100
MARSH REEDGRASS ( <i>Calamagrostis canadensis</i> ) <sup>2</sup>		0-5	67
<b>Moss</b>	<b>59</b>	<b>31-75</b>	<b>100</b>

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1390-1560 M (1451 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
NORTHERLY  
SLOPE:  
0-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 478 (283-672)  
FORBS: 192 (170-214)  
SHRUBS: 252 (204-300)  
TOTAL: 845 (644-1046)

<p>SUGGESTED GRAZING CAPACITY NON-USE</p>
---

## UFE4. Lodgepole pine/ Marsh reedgrass

(*Pinus contorta*/ *Calamagrostis canadensis*)

n=3 This community type is similar to the P1 /hairy wildrye/ fireweed-peavine community type described by Lane et al. (2000). The tree canopy is open which allows good understory growth.

The good understory forage production and easy access through this community type makes it useful for livestock grazing. If this community type occurs adjacent to a physical feature that attracts livestock to the area, it may be considered primary range.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### TREES

LOGEPOLE PINE  
(*Pinus contorta*) 13 0-30 67

WHITE SPRUCE  
(*Picea glauca*) 12 0-20 67

#### SHRUBS

PRICKLY ROSE  
(*Rosa acicularis*) 1 T-2 100

BOG CRANBERRY  
(*Vaccinium vitis-idaea*) 2 0-7 33

BUNCHBERRY  
(*Cornus canadensis*) 5 2-9 100

TWIN-FLOWER  
(*Linnaea borealis*) 4 T-6 100

#### FORBS

FIREWEED  
(*Epilobium angustifolium*) 3 2-3 100

LINDLEY'S ASTER  
(*Aster ciliolatus*) 3 T-6 100

#### GRASSES

HAIRY WILD RYE  
(*Elymus innovatus*) 5 2-6 100

MARSH REEDGRASS  
(*Calamagrostis*

*canadensis*) 12 4-18 100

MOSS 12 8-17 100

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

MESIC

#### NUTRIENT REGIME:

MESOTROPHIC

#### ELEVATION:

1350-1380 M (1367) M

#### SOIL DRAINAGE:

WELL

#### ASPECT:

VARIABLE

#### SLOPE:

0-45%

### FORAGE PRODUCTION(KG/HA)

TOTAL: 801 (600-1200)

SUGGESTED GRAZING CAPACITY

NON-USE

**UFE5. Black spruce/ Willow**  
(*Picea mariana/ Salix spp.*)

n=2 This community type is characterized by a dominant cover of black spruce and a sparse understory cover. The sites are moist in the spring and dry out later in the growing season. Corns and Annas (1986), found that these forests have a fire origin and can persist for more than 150 years.

This community type would be considered non-use for domestic livestock.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

<b>TREES</b>			
BLACK SPRUCE ( <i>Picea mariana</i> )	15	10-20	100
WHITE SPRUCE ( <i>Picea glauca</i> )	6	2-10	100
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	49	32-65	100
LABRADOR TEA ( <i>Ledum groenlandicum</i> )	7	0-14	50
BUNCHBERRY ( <i>Cornus canadensis</i> )	7	0-13	50
<b>FORBS</b>			
PALMATE LEAVED COLTSFOOT ( <i>Petasites palmatus</i> )	1	0-1	50
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	1	T-1	100
HORSETAIL ( <i>Equisetum spp</i> )	8	T-8	100
<b>GRASSES</b>			
WATER SEDGE ( <i>Carex aquatilis</i> )	5	0-10	50
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	4	0-8	50
MOSS	51	41-59	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
HYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1415-1454 M (1435) M  
SOIL DRAINAGE:  
IMPERFECTLY  
ASPECT:  
NORTH  
SLOPE:  
0-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 89 (62-116)  
FORBS: 166 (15-316)  
SHRUBS: 130 (30-230)  
TOTAL: 385 (161-608)

SUGGESTED GRAZING CAPACITY  
NON-USE

**UFE6. White spruce/ Horsetail/ Moss**  
*(Picea glauca/ Equisetum arvense/ Moss spp.)*

n=3 This community type is successionaly more advanced than the Pl-Sw/ bunchberry community type (UFE2) previously described. The lack of fire disturbance has allowed white spruce to succeed into the lodgepole pine canopy and dominate the site. As these stands mature, their canopies close, shading the understory vegetation and allowing moss cover to increase. The sparseness and low palatability of the vegetation limits the use of these stands by domestic livestock.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i>	43	15-65	100
BALSAM POPLAR <i>(Populus balsamifera)</i>	2	0-3	67
<b>UNDERSTORY TREES</b>			
BALSAM POPLAR <i>(Populus balsamifera)</i>	2	0-5	33
<b>SHRUBS</b>			
LOW BUSH CRANBERRY <i>(Viburnum edule)</i>	2	0-3	67
ROSE <i>(Rosa acicularis)</i>	5	0-14	67
TWINFLOWER <i>(Linnaea borealis)</i>	5	0-9	67
<b>FORBS</b>			
PALMATE LEAVED COLTSFOOT <i>(Petasites palmatus)</i>	11	0-22	67
SCOURING RUSH <i>(Equisetum scirpoides)</i>	9	0-16	67
HORSETAIL <i>(Equisetum arvense)</i>	9	9-12	100
TALL LUNGWORT <i>(Mertensia paniculata)</i>	2	0-4	67
<b>GRASSES</b>			
HAIRY WILD RYE <i>(Elymus innovatus)</i>	5	3-7	100
MOSS	37	0-91	67

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1350-1454 M (1415 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 NORTHEAST  
 SLOPE:  
 3%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 83 (68-96)  
 FORBS: 223 (212-234)  
 SHRUBS: 98 (0-196)  
 TOTAL: 418 (332-504)

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFE7. White spruce/ Willow**  
(*Picea glauca/ Salix spp.*)

n=1 This community type is similar to the Sw/ bunchberry/ moss community type, but is found on wetter sites, with poorer drainage. The wetter sites favour the growth of willow in the understory. The high cover of willow and spruce limits the amount of light reaching the understory. Consequently, there is little forage for domestic livestock.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

	MEAN	RANGE	CONST.
<b>TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> )	45	-	100
LOGEPOLE PINE ( <i>Pinus contorta</i> )	10	-	100
<b>SHRUBS</b>			
WILLOW ( <i>Salix spp.</i> )	60	-	100
BOG BIRCH ( <i>Betula glandulosa</i> )	8	-	100
LOW BILBERRY ( <i>Vaccinium caespitosum</i> )	6	-	100
TWIN-FLOWER ( <i>Linnaea borealis</i> )	5	-	100
<b>FORBS</b>			
CREAM COLORED VETCHLING ( <i>Lathyrus ochroleucus</i> )	2	-	100
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	9	-	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	12	-	100
ALPINE ASTER ( <i>Aster alpinus</i> )	3	-	100
YARROW ( <i>Achillea millefolium</i> )	3	-	100
FIREWEED ( <i>Epilobium angustifolium</i> )	3	-	100
<b>GRASSES</b>			
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	7	-	100
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	8	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC  
NUTRIENT REGIME:  
PERMESOTROPHIC  
ELEVATION:  
1646 M  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
WEST  
SLOPE:  
15%

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 300

SUGGESTED GRAZING CAPACITY  
NON-USE

**UFE8. White spruce/ Bearberry**  
*(Picea glauca/ Arctostaphylos uva-ursi)*

n=1 This community type is similar to the Sw/ buffaloberry/ bearberry c.t. described by Lane et al. (2000) in the Lower Foothills. This type is fairly dry with a poor nutrient regime; as indicated by the high abundance of bearberry. It may also be somewhat windswept and desiccated, as indicated by the low tree canopy cover.

If this community type is located near a physical feature that attracts livestock to the area it may be considered to be primary or secondary range. In other instances though, where it is not near an attractive feature, this community type would be considered non-use.

**PLANT COMPOSITION** **CANOPY COVER(%)**

MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i>	20	-	100
ASPEN <i>(Populus tremuloides)</i>	8	-	100
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	12	-	100
WILLOW SPP <i>(Salix spp)</i>	9	-	100
BOG BIRCH <i>(Betula glandulosa)</i>	7	-	100
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	23	-	100
<b>FORBS</b>			
SHOWY LOCOWEED <i>(Oxytropis splendens)</i>	10	-	100
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	18	-	100
ALPINE MILKVETCH <i>(Astragalus alpinus)</i>	7	-	100
CLOVER <i>(Trifolium repens)</i>	6	-	100
DANDELION <i>(Taraxacum officinale)</i>	6	-	100
<b>GRASSES</b>			
PURPLE OATGRASS <i>(Schizachne purpurascens)</i>	18	-	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	14	-	100
BLUNT SEDGE <i>(Carex obtustata)</i>	10	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1311 M  
SOIL DRAINAGE:  
MODERATELY WELL

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 400

<p><b>SUGGESTED GRAZING CAPACITY</b> NON-USE</p>
--

**UFE9. White spruce/ Juniper-Buffaloberry**  
*(Picea glauca/ Juniperus horizontalis-Shepherdia canadensis)*

n=2 This community type was described along the north shore of Brule lake. It is characteristic of the fine-textured, calcareous loess deposits which have blown down the Athabasca river valley from Jasper National Park. The soils of this community have a high pH (8) which supports a good cover of hairy wildrye. This community type is extremely slow growing. When harvested, the cutblocks resemble native grasslands (juniper/ hairy wildrye (UFF1) and rose/ hairy wildrye (UFF2)).

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE			
<i>(Picea glauca)</i> -	50	50	100
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL			
<i>(Potentilla fruticosa)</i>	3	1-4	100
CREEPING JUNIPER.			
<i>(Juniperus horizontalis)</i>	44	43-45	100
PRICKLY ROSE			
<i>(Rosa acicularis)</i>	7	6-8	100
BUFFALOBERRY			
<i>(Shepherdia canadensis)</i>	3	0-5	50
<b>FORBS</b>			
WHITE CAMAS			
<i>(Zigadenus elegans)</i>	4	T-7	100
SHOWY LOCOWEED			
<i>(Oxytropis splendens)</i>	2	T-3	100
NORTHERN HEDYSARUM			
<i>(Hedysarum boreale)</i>	2	0-4	50
NORTHERN BEDSTRAW			
<i>(Galium boreale)</i>	2	1-2	100
BASTARD TOAD FLAX			
<i>(Comandra umbellata)</i>	1	1-2	100
<b>GRASSES</b>			
HAIRY WILDRYE			
<i>(Elymus innovatus)</i>	14	13-14	100
BLUNT SEDGE			
<i>(Carex obtustata)</i>	4	3-5	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1066 M  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 SOUTH  
 SLOPE:  
 0-5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 297 (294-300)  
 FORBS: 176 (146-206)  
 SHRUBS: 181 (36-326)  
 TOTAL: 654 (536-772)

SUGGESTED GRAZING CAPACITY  
 NON-USE

**UFE10. White spruce/ Moss**  
(*Picea glauca*/ Moss spp.)

n=1 This community type represents a successional mature forested stand in the Upper Foothills subregion. As succession occurs from pine to spruce, the canopy cover becomes closed and the amount of understory vegetation decreases until most of the shrub, forb and grass layers have been eliminated. As a result, there is limited forage available for domestic livestock within these spruce dominated community types. This community is typically rated as non-use for domestic livestock.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> ) -	45	-	100
LODGEPOLE PINE ( <i>Pinus contorta</i> )	5	-	100
<b>UNDERSTORY TREES</b>			
SUBALPINE FIR ( <i>Abies lasiocarpa</i> )	25	-	100
<b>SHRUBS</b>			
BUFFALOBERRY ( <i>Shepherdia canadensis</i> )	5	-	100
BOG CRANBERRY. ( <i>Vaccinium caespitosum</i> )	4	-	100
PRICKLY ROSE ( <i>Rosa acicularis</i> )	2	-	100
TWINFLOWER ( <i>Linnaea borealis</i> )	10	-	100
<b>FORBS</b>			
HEART LEAVED ARNICA ( <i>Arnica cordifolia</i> )	8	-	100
SHOWY ASTER ( <i>Aster conspicuus</i> )	5	-	100
BUNCHBERRY ( <i>Cornus canadensis</i> )	3	-	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	2	-	100
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	10	-	100
<b>MOSS</b>			
STAIR STEP MOSS ( <i>Hylocomium splendens</i> )	90	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1350 M  
SOIL DRAINAGE:  
WELL  
ASPECT:  
EAST  
SLOPE:  
10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 78  
FORBS: 96  
SHRUBS: 160  
TOTAL: 332

SUGGESTED GRAZING CAPACITY  
NON-USE

[Faint, illegible text at the top of the page, possibly a header or title area.]

[Large block of very faint, illegible text, likely the main body of the document.]

[Faint, illegible text at the bottom of the page, possibly a footer or concluding remarks.]

## Cutblock and burn community types

In general, cutblocks provide only a limited source of forage for domestic livestock in the Upper Foothills subregion. The Brule stock association, Robb head tax permit and Upper James allotment are examples where the livestock rely principally on the forage within harvested cutblocks. On average, cutblocks produce twice as much forage as deciduous stands and nearly 3 times the forage as coniferous stands. In the Brule stock association, forage production on the cutblocks averaged 3-5 times greater than the unharvested white spruce dominated forest.

Two of the cutblock community types in this guide were described from the Brule stock association. These are the juniper/ hairy wildrye (UFF1) and rose/ hairy wildrye (UFF2) c.t.. Both of these types have very little growth of regenerating trees and resemble native grasslands (Figure 7).


Other cutblock community types were described on moister sites throughout the Solomon valley. These communities represent areas that were harvested 30-40 years ago. Currently, they are important sources of forage for domestic livestock throughout the area.

One burn community type was described from the Solomon valley. This burn occurred on an Se-Fa/ willow community approximately 10 years ago. The site was located in an area that had nutrient rich seepage that made it very productive for horses grazing the area.

In the Upper James and Wilson creek allotments west of Sundre, harvesting of lodgepole pine dominated sites produces fireweed/ hairy wildrye dominated communities on south and west facing slopes. On the more northern aspects in this area, the cutblocks tended to be dominated by moss to form the Pl-Sw/ moss community type. Livestock preferred to graze the fireweed/ hairy wildrye dominated cutblocks.



**Figure 7. The juniper/ hairy wildrye community type (UFF1) results from the harvesting of a Sw/ juniper community.**



**UFF1. Juniper/ Hairy wildrye**  
(*Juniperus horizontalis/ Elymus innovatus*)

n=4 This community represents a harvested Sw/ juniper community along the north shore of Brule lake. It is characteristic of the fine-textured, calcareous loess deposits, which have blown down the Athabasca river valley from Jasper National Park. The soils of this community have a high pH (8) which supports a good cover of hairy wildrye. This community type is extremely slow growing. When harvested, the cutblocks resemble native grasslands.

This community is very similar to the rose/ hairy wildrye community, but appears to be in a later successional stage. This community type was described in older cutblocks (35 yrs) than the rose/ hairy wildrye community type (UFF2). As succession occurs on these cutblocks it appears that juniper and grass cover increase, causing a corresponding increase in forage production.

**PLANT COMPOSITION CANOPY COVER(%)**

	Mean	Range	Const.
<b>TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> )	12	5-18	100
BALSAM POPLAR ( <i>Populus balsamifera</i> )	8	0-15	75
ASPEN ( <i>Populus tremuloides</i> )	6	0-15	50
<b>UNDERSTORY TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> )	1	0-2	25
BALSAM POPLAR ( <i>Populus balsamifera</i> )	1	0-1	25
ASPEN ( <i>Populus tremuloides</i> )	1	0-2	50
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	7	2-11	100
CREEPING JUNIPER ( <i>Juniperus horizontalis</i> )	19	11-27	100
PRICKLY ROSE ( <i>Rosa acicularis</i> )	6	0-10	75
WILLOW ( <i>Salix spp.</i> )	14	3-15	100
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	9	0-17	75
<b>FORBS</b>			
SHOWY LOCOWEED ( <i>Oxytropis splendens</i> )	3	1-4	100
NORTHERN HEDYSARUM ( <i>Hedysarum boreale</i> )	6	0-7	75
NORTHERN BEDSTRAW ( <i>Galium boreale</i> )	8	6-10	100
<b>GRASSES</b>			

HAIRY WILDRYE ( <i>Elymus innovatus</i> )	12	3-24	100
BLUNT SEDGE ( <i>Carex obtusata</i> )	6	0-15	75
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> ) <sup>2</sup>		0-4	50

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME: SUBMESIC
NUTRIENT REGIME: SUBMESOTROPHIC
ELEVATION: 1036-1066 M (1046 M)
SOIL DRAINAGE: WELL
ASPECT: SOUTH
SLOPE: 0-5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 520 (268-866)
FORBS: 697 (124-1538)
SHRUBS: 267 (12-450)
TOTAL: 2089 (592-3732)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM OR 1.0 AC/AUM

**UFF2. Rose/ Hairy wildrye**  
(*Rosa acicularis/ Elymus innovatus*)

n=10 This community type represents a Sw/ juniper community that was harvested 20 years ago. It is very similar to the previously described juniper/ hairy wildrye community (UFF1), but lacks the cover of juniper. It appears that harvesting disturbance causes juniper to decline in cover. As succession occurs, juniper and grass density increase, causing forage productivity to increase. The site conditions are so harsh it appears that grass cover has to undergo succession onto the site.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

**TREES**

WHITE SPRUCE  
(*Picea glauca*)- 9 0-20 90

BALSAM POPLAR  
(*Populus balsamifera*) 6 0-15 80

ASPEN  
(*Populus tremuloides*) 2 0-10 30

**UNDERSTORY TREES**

WHITE SPRUCE  
(*Picea glauca*) 2 0-15 10

BALSAM POPLAR  
(*Populus balsamifera*) 2 0-20 50

ASPEN  
(*Populus tremuloides*) 1 0-1 20

**SHRUBS**

SHRUBBY CINQUEFOIL  
(*Potentilla fruticosa*) 1 0-4 80

CREEPING JUNIPER.  
(*Juniperus horizontalis*) 2 0-7 60

PRICKLY ROSE  
(*Rosa acicularis*) 4 0-13 90

WILLOW SPP.  
(*Salix spp.*) 6 0-10 80

BEARBERRY  
(*Arctostaphylos uva-ursi*) 1 0-7 50

**FORBS**

WHITE CAMAS  
(*Zigadenus elegans*) 1 0-3 30

SHOWY LOCOWEED  
(*Oxytropis splendens*) 2 0-4 60

NORTHERN HEDYSARUM  
(*Hedysarum boreale*) 1 0-24 40

NORTHERN BEDSTRAW  
(*Galium boreale*) 4 T-11 100

DANDELION  
(*Taraxacum officinale*) 3 0-8 90

**GRASSES**

HAIRY WILDRYE  
(*Elymus innovatus*) 24 4-40 100  
BLUNT SEDGE  
(*Carex obtustata*) 2 0-7 60  
SLENDER WHEATGRASS  
(*Agropyron trachycaulum*)4 0-10 70

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1036 M

SOIL DRAINAGE:

WELL

ASPECT:

SOUTHERLY

SLOPE:

2-10%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 723 (212-1514)

FORBS: 388 (126-756)

SHRUBS: 132 (2-454)

TOTAL: 1243 (540-2360)

**SUGGESTED GRAZING CAPACITY**

0.7 HA/AUM OR 1.6 AC/AUM

**UFF2a. Fireweed/ Hairy wildrye**  
*(Epilobium angustifolium/ Elymus innovatus)*

n=22 This community type represents a P1/ moss community that was harvested 5-7 years ago. This community type was described on south and west facing slopes throughout the area. On more northerly aspects, moss dominates the understory of these cutblocks. Cutblocks can be an important source of forage for domestic livestock. They produce on average twice as much as deciduous stands, and nearly three times more than conifer stands. It must be remembered that this increase in forage is only temporary. As the cutblock undergoes succession there is a corresponding drop in production.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>UNDERSTORY TREES</b>			
LOGEPOLE PINE <i>(Pinus contorta)</i>	1	0-10	59
ASPEN <i>(Populus tremuloides)</i>	1	0-2	23
<b>SHRUBS</b>			
PRICKLY ROSE <i>(Rosa acicularis)</i>	1	0-5	86
WILLOW SPP. <i>(Salix spp.)</i>	1	0-6	41
BUNCHBERRY <i>(Cornus canadensis)</i>	1	0-1	73
<b>FORBS</b>			
FIREWEED <i>(Epilobium angustifolium)</i>	5	0-7	91
SHOWY ASTER <i>(Aster conspicuus)</i>	1	0-7	46
NORTHERN BEDSTRAW <i>(Galium boreale)</i>	1	0-1	27
<b>GRASSES</b>			
HAIRY WILDRYE <i>(Elymus innovatus)</i>	12	0-16	96
SEDGE <i>(Carex spp)</i>	3	0-9	91
PINEGRASS <i>(Calamagrostis rubescens)</i>	1	0-11	27

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBMESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1390-1700 M (1513 M)  
 SOIL DRAINAGE:  
 WELL  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 2-30%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 704 (566-842)  
 FORBS: 511 (178-844)  
 TOTAL: 1215 (744-1686)

**SUGGESTED GRAZING CAPACITY**  
 0.7 HA/AUM OR 1.7 AC/AUM

**UFF3. White spruce/ Horsetail/ Kentucky bluegrass**  
*(Picea glauca/ Equisetum arvense/ Poa pratensis)*

n=1 This community type represents a Sw/ Horsetail community that was harvested 30-40 years ago along the banks of Moosehorn creek east of Rock Lake. These cutblocks are an important source of forage for domestic livestock and have been extensively utilized by cattle throughout the summer months. The high moisture and nutrient content of the sites make them extremely productive. Once invaded by agronomic species (Kentucky bluegrass and clover) they are extremely palatable to livestock.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i> -	40	-	100
<b>SHRUBS</b>			
WILLOW <i>(Salix spp.)</i>	2	-	100
PRICKLY ROSE <i>(Rosa acicularis)</i>	3	-	100
<b>FORBS</b>			
DANDELION <i>(Taraxacum officinale)</i>	5	-	100
TALL LARKSPUR <i>(Delphinium glaucum)</i>	5	-	100
DEWBERRY <i>(Rubus pubescens)</i>	5	-	100
TALL LUNGWORT <i>(Mertensia paniculata)</i>	5	-	100
YARROW <i>(Achillea millefolium)</i>	3	-	100
HORSETAIL <i>(Equisetum arvense)</i>	2	-	100
<b>GRASSES</b>			
HAIRY WILDRYE <i>(Elymus innovatus)</i>	3	-	100
KENTUCKY BLUEGRASS <i>(Poa pratensis)</i>	12	-	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	3	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBHYGRIC  
 NUTRIENT REGIME:  
 PERMESOTROPHIC  
 ELEVATION:  
 1350 M  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 NORTH  
 SLOPE:  
 1%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 498  
 FORBS: 2378  
 TOTAL: 2876

**SUGGESTED GRAZING CAPACITY**  
 0.3 HA/AUM OR 0.7 AC/AUM

## UFF4. White spruce/ Moss

(*Picea glauca*/ Moss spp.)

n=1 This community type represents a Sw/ moss community that was harvested 30-40 years ago along the banks of West Solomon Creek. The regeneration on this cutblock is to subalpine fir which is similar to the understory of the Sw/ moss (UFE10) community that was harvested in the same area. This community is an important source of forage for wintering horses. The open canopy cover allows for a greater abundance of forbs and grasses in the understory. As the community continues to undergo succession and the canopy becomes denser there will be a corresponding drop in available forage.

### PLANT COMPOSITION CANOPY COVER(%) MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE ( <i>Picea glauca</i> )	10	-	100
SUBALPINE FIR ( <i>Abies lasiocarpa</i> )	30	-	100
<b>SHRUBS</b>			
WILLOW ( <i>Salix spp.</i> )	3	-	100
RIVER ALDER ( <i>Alnus tenuifolia</i> )	3	-	100
BUNCHBERRY ( <i>Cornus canadensis</i> )	1	-	100
<b>FORBS</b>			
FIREWEED ( <i>Epilobium angustifolium</i> )	4	-	100
PEAVINE ( <i>Lathyrus ochroleucus</i> )	1	-	100
<b>MOSS</b>			
FEATHER MOSS ( <i>Pleurozium scherberi</i> )	6	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME:  
MESOTROPHIC  
ELEVATION:  
1300 M  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
EAST  
SLOPE:  
10%

### FORAGE PRODUCTION(KG/HA)

GRASS: 428  
FORBS: 476  
SHRUBS: 78  
TOTAL: 982

SUGGESTED GRAZING CAPACITY  
0.9 HA/AUM OR 2.1 AC/AUM

**UFF4a. Lodgepole pine-White spruce/ Moss**  
*(Pinus contorta-Picea glauca/ Moss spp.)*

n=6 This community type represents a Sw/ moss or Lodgepole pine community that was harvested 5-10 years ago. These moss dominated cutblocks tend to occupy north aspects where the climatic conditions are cooler and moister. Livestock do not prefer to graze these sites.

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

<b>TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i>	3	0-15	17
LODGEPOLE PINE <i>(Pinus contorta)</i>	10	0-25	50
<b>UNDERSTORY TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i>	2	0-8	50
LODGEPOLE PINE <i>(Pinus contorta)</i>	2	0-10	17
<b>SHRUBS</b>			
WILLOW <i>(Salix spp.)</i>	1	0-2	67
ROSE <i>(Rosa acicularis)</i>	1	0-2	100
BUNCHBERRY <i>(Cornus canadensis)</i>	2	0-10	50
<b>FORBS</b>			
FIREWEED <i>(Epilobium angustifolium)</i>	1	0-5	50
HORSETAIL <i>(Equisetum arvense)</i>	5	0-18	50
<b>GRASS</b>			
HAIRY WILDRYE <i>(Elymus innovatus)</i>	6	0-13	100
MARSH REEDGRASS <i>(Calamagrostis canadensis)</i>	3	0-8	33
<b>MOSS</b>			
FEATHER MOSS <i>(Pleurozium scherberi)</i>	1	0-3	50
STAIR STEP MOSS <i>(Hylocomium splendens)</i>	6	0-15	67

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC  
 NUTRIENT REGIME:  
 MESOTROPHIC  
 ELEVATION:  
 1335-1572 M (1414 M)  
 SOIL DRAINAGE:  
 MODERATELY WELL  
 ASPECT:  
 VARIABLE  
 SLOPE:  
 0-16%

**FORAGE PRODUCTION(KG/HA)**

TOTAL: 450

SUGGESTED GRAZING CAPACITY  
 NON-USE



**UFF6. Aspen/ Fireweed**  
(*Populus tremuloides/ Epilobium angustifolium*)

n=1 This community type represents a Pl-Sw/ bunchberry community that was harvested near the Robb area. The regeneration of this community back to aspen indicates that this particular cutblock is transitional to the Lower Foothills subregion. Indeed, the Robb area is on the border between the Upper and Lower Foothills subregions. This community type is highly productive for domestic livestock. Harvesting the trees allows the grasses and forbs to grow, increasing the forage productivity.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

<b>UNDERSTORY TREES</b>			
WHITE SPRUCE			
( <i>Picea glauca</i> )-	1	-	100
ASPEN			
( <i>Populus tremuloides</i> )	6	-	100
<b>SHRUBS</b>			
PRICKLY ROSE.			
( <i>Rosa acicularis</i> )	6	-	100
GREEN ALDER			
( <i>Alnus crispa</i> )	7	-	100
DEWBERRY			
( <i>Rubus pubescens</i> )	3	-	100
<b>FORBS</b>			
FIREWEED			
( <i>Epilobium angustifolium</i> )	52	-	100
HORSETAIL			
( <i>Equisetum arvense</i> )	9	-	100
HEART LEAVED ARNICA			
( <i>Arnica cordifolia</i> )	8	-	100
TALL LUNGWORT			
( <i>Mertensia paniculata</i> )	2	-	100
<b>GRASSES</b>			
SEDGE spp.			
( <i>Carex sp.</i> )	3	-	100
MARSH REEDGRASS			
( <i>Calamagrostis canadensis</i> )	13	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC  
NUTRIENT REGIME :  
PERMESOTROPHIC  
ELEVATION:  
1091 M  
SOIL DRAINAGE :  
MODERATELY WELL  
ASPECT:  
NORTH  
SLOPE:  
5%

**FORAGE PRODUCTION(KG/HA)**

GRASS: 540  
FORBS: 1520  
SHRUBS: 150  
TOTAL: 2210

SUGGESTED GRAZING CAPACITY  
0.8 HA/AUM OR 1.8 AC/AUM

## UFF7. Aspen/ Blueberry-Bearberry/ Hairy wildrye

(*Populus tremuloides*/ *Vaccinium myrtilloides*-*Arctostaphylos uva-ursi*/ *Elymus innovatus*)

n=2 This community type represents a P1/ bog cranberry community (UFE1) that was harvested near the Robb area. The ecological conditions of this site are drier with a poorer nutrient regime. Regeneration of the trees will be much slower than the Aw/ fireweed community type which was described previously. Harvesting of the pine overstory allows grasses and forbs to flourish. This provides a good forage base for domestic livestock. Caution must be used when grazing cutblocks that the stocking rate is not too high to limit the growth of regenerating trees.

### PLANT COMPOSITION CANOPY COVER(%) MEAN RANGE CONST.

#### UNDERSTORY TREES

WHITE SPRUCE

(*Picea glauca*) 1 0-1 50

ASPEN

(*Populus tremuloides*) 7 0-13 50

LODGEPOLE PINE

(*Pinus contorta*) 2 T-3 100

#### SHRUBS

BLUEBERRY

(*Vaccinium myrtilloides*) 11 T-22 100

BOG CRANBERRY

(*Vaccinium caespitosum*) 2 0-3 50

#### FORBS

FIREWEED

(*Epilobium angustifolium*) 1 0-2 50

LINDLEY'S ASTER

(*Aster ciliolatus*) 2 0-4 50

STRAWBERRY

(*Fragaria virginiana*) 2 T-3 100

YARROW

(*Achillea millefolium*) 1 0-2 50

#### GRASSES

INDIAN RICEGRASS

(*Oryzopsis pungens*) 2 T-3 100

HAIRY WILDRYE

(*Elymus innovatus*) 6 T-10 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :

SUBMESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1091 M

SOIL DRAINAGE :

WELL

ASPECT:

EAST

SLOPE:

3%

### FORAGE PRODUCTION(KG/HA)

GRASS: 300

FORBS: 310

SHRUBS: 285

TOTAL: 895

SUGGESTED GRAZING CAPACITY

2.0 HA/AUM OR 4.6 AC/AUM

## UFF8. Kentucky bluegrass-Creeping red fescue/ Clover

(*Poa pratensis*-*Festuca rubra*/ *Trifolium spp.*)

n=5 This community type represents cutblocks that have been heavily grazed by livestock. Heavy livestock grazing favours the growth of the invaders Kentucky bluegrass and timothy. The grazing pressure which favours the growth of these grass species is usually detrimental to the growth of trees. Cattle damage to the conifer trees is usually trampling damage which scars the trees and breaks the stem.

### PLANT COMPOSITION CANOPY COVER(%) MEAN RANGE CONST.

<b>SHRUBS</b>			
ROSE			
( <i>Rosa acicularis</i> )	1	0-4	20
<b>FORBS</b>			
<b>FIREWEED</b>			
( <i>Epilobium angustifolium</i> )	1	0-3	80
<b>CLOVER</b>			
( <i>Trifolium spp</i> )	15	0-48	80
<b>STRAWBERRY</b>			
( <i>Fragaria virginiana</i> )	1	0-2	40
<b>YARROW</b>			
( <i>Achillea millefolium</i> )	1	0-T	60
<b>GRASSES</b>			
<b>TIMOTHY</b>			
( <i>Phleum pratense</i> )	8	T-35	100
<b>HAIRY WILDRIE</b>			
( <i>Elymus innovatus</i> )	1	0-3	40
<b>KENTUCKY BLUEGRASS</b>			
( <i>Poa pratensis</i> )	15	0-67	80
<b>CREEPING RED FESCUE</b>			
( <i>Festuca rubra</i> )	10	0-41	80

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBMESIC  
NUTRIENT REGIME:  
SUBMESOTROPHIC  
ELEVATION:  
1435-1480 M (1453 M)  
SOIL DRAINAGE:  
MODERATELY WELL  
ASPECT:  
VARIABLE  
SLOPE:  
0-12%

### FORAGE PRODUCTION(KG/HA)

GRASS: 932  
TOTAL: 932

SUGGESTED GRAZING CAPACITY  
2.0 HA/AUM OR 4.6 AC/AUM

### Literature cited

- Bailey, A.W., M.G. Willoughby, R. Johansen and S. Smith. Management of Yukon Rangelands. Renewable Resources, Yukon Territorial Government, Whitehorse, Yukon. 55pp. ISBN-1-55018-138-6.
- Bork, E. 1990. Clearwater allotment: Prescribed burn vegetation survey, evaluation, and discussion. Range Management Section, Alberta Forest Service, Edmonton, Alta. 29pp.
- Bork, E. 1994. Ecological classification and management of native ranges in Willmore Wilderness Park. Alberta Environmental Protection. Forest Management Division. Edmonton, Alta. Pub. no. T/282. 65pp.
- Beckingham, J. 1994. Field guide to the ecosystems of Northern Alberta. Environmental Protection, Land and Forest Services, Edmonton, Alta. 352pp.
- Beckingham, J.D., I.G.W. Corns and J.H. Archibald. 1996. Field guide to ecosites of West-Central Alberta. Nat. Resour. Can., Can. For. Serv., Northwest Reg., North. For. Cent., Edmonton, Alberta. Spec. Rep. 9.
- Corns, I.G.W. and P.L. Achuff. 1982. Vegetation type descriptions. In Holland, W.D. and G.M. Coen. Ecological (Biophysical) Land Classification of Banff and Jasper National Parks. Vol II. Soil and Vegetation Resources. Environment Canada and Alberta Institute of Pedology, University of Alberta. Pub. no. SS-82-44. 540pp.
- Corns, I.G.W. and R.M. Annas. 1986. Field guide to forest ecosystems of West-Central Alberta. Northern Forestry Center, Canadian Forestry Service, Edmonton, Alta. 251pp.
- Daubenmire, R. 1952. Forest vegetation of Northern Idaho and adjacent Washington and its bearing on concepts of vegetation classification. *Ecol. Mongr.* 22: 301-330.
- Daubenmire, R. 1978. *Plant Geography*. Academic Press. New York. 338pp.
- Dept. of Environmental Protection. 1994. Natural regions of Alberta. Alberta Environmental Protection. Edmonton, Alta. Pub. no. I/531. 18pp.
- Gauch, H.G. 1982. *Multivariate analysis in community ecology*. Cambridge University Press, Cambridge, 298pp.
- Lane, C. T., M.G. Willoughby and M.J. Alexander. 2000. Range plant community types and carrying capacity for the Lower Foothills subregion of Alberta. Alberta Environment and Alberta Agriculture Food and Rural Development. Edmonton. AB. Pub. no. T/532. 232pp.

Mackinnon, A., J. Pojar, and R. Coupe. 1992. *Plants of Northern British Columbia*. Lone Pine Publishing, Edmonton, Alta. 345pp.

Mueggler, W.F. 1988. *Aspen community types of the Intermountain Region*. U.S.D.A. Intermountain Research Station. INT-250. 133pp.

Rangeland Resource Information System. 1991. *Outline and guide, a proposal*. Range Management Section, Alberta Forest Service, Edmonton, Alta. 19pp.

Range Survey Manual. 1992. Range Management Section, Alberta Forest Service. Edmonton, Alta. 39pp.

Strong, W.L. and H.G. Anderson. 1980. *Ecological land classification and evaluation reference manual*. Resource Evaluation reference manual. Resource Evaluation Branch, Energy and Natural Resources. Edmonton, Alta. 160pp.

Strong, W.L. and K.R. Leggat. 1992. *Ecoregions of Alberta*. Alberta Forestry, Lands and Wildlife, Resource Information Branch, Edmonton, Alta. T/245. 77pp.

Strong, W.L. 1992. *Ecoregions and Ecodistricts of Alberta*. Alberta Forestry, Lands and Wildlife. Edmonton, Alta. Pub. no. T/244. 77pp.

Willoughby, M.G. 1992. *Rangeland Reference Areas, Plant communities, ecology and response to grazing in Division 3*. Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alta. T/269, 36pp.

Willoughby, M.G. 1995. *The effect of grazing on deciduous plant communities in the Boreal Ecoprovince of Alberta*. Fifth International Rangeland Congress. Salt Lake City Utah. pg. 608-609.

Willoughby, M.G. 1999. *Range plant community types and carrying capacity for the Subalpine and Alpine subregions of Alberta*. Environmental Protection. Land and Forest Services. Edmonton, Alta. Pub. no. T/438. 109pp.

Willoughby, M.G. 2000. *Guide to range plant communities and carrying capacity for the Dry and Central Mixedwood subregions of Alberta*. Alberta Environment and Alberta Agriculture Food and Rural Development. Edmonton. AB. Pub. no. T/526. 187pp.

Willoughby, M.G. 2000. *Rangeland Reference Areas: Species composition changes in the presence and absence of grazing and fire on the Rough fescue-Hairy wildrye dominated community types of the Upper Foothills subregion*. Alberta Environment. Edmonton. AB. Pub. no. T/572. 30pp.

Willoughby, M.G., M.J. Alexander and K.M. Sundquist. 2001. Range plant community types and carrying capacity for the Montane subregion. Alberta Environmental Protection. Edmonton. AB. Pub. no. T/589. 194 pp.

Youngblood, A. Community type classification of forest vegetation in young, mixed stands, Interior Alaska. U.S.D.A. Forest Service, Pacific Northwest Research Station. PNW-RP-458. 42pp.

Willoughby, M.G. 1982. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.

Willoughby, M.G. 1984. A guide to range plant communities and carrying capacity for the Dry and Central Mixedwood Subregions of Alberta. Alberta Environment and Alberta Agriculture Food and Rural Development, Edmonton, Alberta. Pub. no. T376. 187 pp.

Willoughby, M.G. 1985. The effect of grazing on deciduous plant communities in the Boreal Ecoregion of Alberta. Fifth International Rangeland Congress, Salt Lake City, Utah. pp. 608-609.

Willoughby, M.G. 1986. Rangeland Reference Areas: Species composition changes in the presence and absence of grazing and fire on the Rough Prairie-Hairy willow dominated community types of the Upper Foothills subregion. Alberta Environment, Edmonton, Alberta. Pub. no. T372. 30 pp.

Willoughby, M.G. 1989. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.

Willoughby, M.G. 1991. Ecotones of Alberta. Alberta Forestry, Lands and Wildlife, Resource Information Branch, Edmonton, Alberta. T045. 77 pp.

Willoughby, M.G. 1992. Ecotones and Ecotrends of Alberta. Alberta Forestry, Lands and Wildlife, Edmonton, Alberta. Pub. no. T244. 77 pp.

Willoughby, M.G. 1993. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.

Willoughby, M.G. 1994. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.

Willoughby, M.G. 1995. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.

Willoughby, M.G. 1996. Rangeland Reference Areas: Plant communities, ecology and response to grazing in Division 3, Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alberta. Pub. no. T282. 194 pp.



National Library of Canada  
Bibliothèque nationale du Canada



3 3286 52476647 0